



Systemic Enzymes and their Role in the Reduction of Post-Operative Edema After Surgical Removal of Lower Third Molars: A Randomized Controlled Double-Blind Study

Abhinav R P¹, M.P. Santhosh kumar^{2*}

¹Associate Professor, Department of Implantology, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai 77

²Professor, Department of Oral and Maxillofacial Surgery, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai 77

***Corresponding author:** M.P. Santhosh kumar, Professor, Department of Oral and Maxillofacial Surgery, Saveetha Dental College and Hospital, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, 162, Poonamallee High Road, Velappanchavadi, Chennai 600077 Tamil Nadu, India, Email: santhoshsurgeon@gmail.com

Submitted: 20 February 2023; Accepted: 23 March 2023; Published: 02 April 2023

ABSTRACT

The most common postoperative complications after surgical removal of impacted mandibular third molar tooth are edema, pain and trismus. Different types of analgesics and anti-inflammatory agents have been employed for reduction of these complications. The aim of the study was to evaluate the effect of systemic enzymes to control post-operative edema, pain and trismus after surgical removal of impacted lower third molars.

The enzyme combination of trypsin, bromelain, rutoside trihydrate was compared with that of serratiopeptidase to control post-operative sequelae following lower third molar surgery. Sixty-eight patients, who required surgical removal of impacted mandibular third molars participated in this study. They were randomly allocated into two groups (n=34 in each group). Group I received 5mg serratiopeptidase orally along with the standard medication and the other group received the enzyme combination with the standard medication. P value <0.05 was considered to be statistically significant, Five patients were lost to follow-up. The mean swelling measures were found to be non-significant when compared between the two groups (Group I vs. II-54.7 ± 3.4 Vs. 53.2 ± 3.2, p=0.34). Similarly, the differences in mean pain and mouth opening values between the study groups was found to be non-significant (p>0.05).

There was no significant difference in the effect of the enzyme combination of rutoside, trypsin and bromelain when compared to serratiopeptidase in the reduction of post-operative edema, pain and trismus following lower third molar surgery.

Keywords: Systemic enzymes, third molar surgery, post-operative complications, pain, edema

INTRODUCTION

The presence of impacted teeth in the mandible can cause problems to the adjacent teeth and mucosa if they do not erupt into the proper

position. A study done by Srivastava N et al concluded that in 55% of the cases, impacted mandibular molars cause distal caries in the second molars [1].

It can also lead to pericoronitis which may proceed to become a space infection involving the pterygomandibular space and considering all of the above issues with an impacted third molar, it is generally recommended to be extracted.

Halpern et al., grouped the postoperative morbidities into complications and immediate postoperative tissue reactions [2]. The immediate postoperative tissue reactions are characterized by pain, swelling, reduced masticatory capability, trismus and dysphagia. The complications are unpredicted reactions that may or may not occur in the immediate post-operative time and they include dry socket bleeding, nerve injury, delayed healing, and infection periodontal pocket. The main goal should be to prevent/reduce these post-operative issues, and a variety of medications and modalities, including physical therapeutic treatments such as cryotherapy and soft laser application, are available to help with this [3,4].

In the existing literature, the medical management of post-operative edema is confined mainly to the use of non-steroidal anti-inflammatory drugs (NSAIDs) or steroids. The use of enzyme treatment including trypsin, bromelain, and rutoside trihydrate for edema control in third molar surgery is not well established. Antioxidant, anti-inflammatory, analgesic, antiplatelet, and immunological modifying effects are all present in this combination [5]. In several clinical studies combination of trypsin, bromelain and rutoside has shown to have significant anti-inflammatory effects [6]. To explore more about this combination, the present study was designed to evaluate the efficacy of systemic enzymes to control postoperative edema after surgical removal of impacted mandibular third molars.

MATERIALS AND METHODS

This study was performed in the Oral and Maxillofacial Surgery Department of Saveetha Dental College and Hospitals, Chennai. Sixty-eight patients who required surgical removal of impacted mandibular third molars were included in the study. They were randomly divided into two groups (Group I- Serratiopeptidase, Group

II- Enzyme combination) of thirty-four subjects each using a random allocation software [7]. This study was reviewed and approved by the Institutional Review Board and informed consent was obtained from all the participants. The inclusion criteria were patients who were healthy individuals, aged 18 years and above, only third molars which were graded as Grade II and above in the Parant's difficulty scale, patients who had bilateral impacted teeth for whom only one side was removed surgically were included [8]. The patients with a positive medical history, those who had consumed analgesics within 24 hours prior to surgery, pregnant or lactating women were excluded from this study.

Group I received 5mg serratiopeptidase orally along with the standard medication (500mg amoxicillin and 500mg paracetamol thrice daily) whereas Group II received the systemic enzyme combination with the standard medication. The proteolytic enzyme combinations used were rutoside, trypsin and bromelain. These three agents were given in the form of an enteric-coated tablet (this to prevent the enzymes from being disintegrated by gastric secretions) that contained 100 mg of rutoside, 48 mg of trypsin, 90 mg of bromelain. Rescue medication, where necessary, was given in the form of ketorolac tablets 10 mg.

The radiographs (orthopantomograph or intraoral periapical radiographs) were taken to assess third molar impaction regarding difficulty using WAR (White, Amber and Red) lines and the Parant scale. In this double blinded prospective randomized clinical trial, the primary outcome variable was postoperative edema which was assessed by measurements made between the 6 reference points of Tragus, corner of the mouth, chin, angle of mandible, external canthus and nasal bone. The secondary outcome variables were mouth opening measured using a calliper and pain assessed using a visual analogue scale. These parameters were recorded on the 1st, 3rd and 7th postoperative days respectively.

Statistical Analysis

The collected results underwent normality tests such the Kolmogrov-Smirnov and Shapiro Wilk's

test and the results revealed that the data followed a parametric distribution. For parametric continuous data, student t-test was performed and to define statistical significance, significance level of $p < 0.05$ was chosen. Statistical software (SPSS v.17 software, IBM, Armonk, New York) was used to conduct all the analyses.

RESULTS

Of a total sixty-eight patients in this study, five patients were later excluded as they were lost during follow up. Hence, a total of 63 patients were included in the final analysis who had attended the follow-up visits. The mean patient age in groups I and II were 27 and 28 years respectively. There were no statistically significant differences in the demographic characteristics of the subjects or the parameters related to the surgical procedure among the study groups.

The mean preoperative edema in both the control and test groups were comparable. The mean postoperative swelling on the 1st postoperative day in Group I was 54.7 mm and in Group II was 53.2 mm which was not statistically significant ($p = 0.34$). There was an increasing trend in facial swelling till the 3rd postoperative day, which decreased on the 7th postoperative day. The highest was observed in both the groups on the third postoperative day and least on the 7th postoperative day. Similarly, the difference between mean facial swellings was highest on the 3rd postoperative day. The mean swelling measures were found to be non-significant when compared between the two groups but a significant difference in mean facial swelling was observed on day 1, day 3 and day 7 compared to the preoperative day within the group ($p < 0.001$) [Table 1].

The mean postoperative pain score immediately after surgery was 8.3 ± 0.9 in group I and 8.2 ± 0.5 in group II. The mean pain score was highest on the day of surgery, which progressively decreased by the 1st, 3rd and 7th postoperative day. The mean pain scores were non-significant when compared between the groups but were significant when compared within the groups ($p < 0.001$) [Table 2].

Analysis of the data showed that the mean baseline measure of the inter-incisal distance was 44 ± 3.7 mm and 43.4 ± 8.2 mm in group I and II respectively. In both the groups, trismus was greater on day one after surgery and had subsided during the subsequent follow-up intervals. There was no significant difference between the Serrapeptase group and the Enzyme combination group in the mean maximal interincisal distance throughout the follow-up period [Table 3]. Data regarding the use of rescue medication in the treatment group is presented in Figure 1. Of the 31 patients recruited in group I, only 3 (14.3%) required rescue medication (ketorolac) as compared to 6 out of 32 patients (18.8%) in group II. As per the Parant scale for difficulty grading for third molar removal, the majority of the cases were classified as a grade 3 difficulty (76.9%).

DISCUSSION

The current study compared the efficacy of an enzyme combination (Trypsin, Bromelain, and Rutoside) with serratiopeptidase on the postoperative sequelae after surgical removal of impacted mandibular third molars. Age, gender, operative time, and surgical experience are all risk factors for edema, pain, and trismus after third molar surgery [9].

Mild bleeding, edema, stiffness, and pain are all common physiologic reactions following the surgical removal of an impacted third molar. All of these are perceived as unpleasant by the patient and should be avoided to the greatest extent possible. The postoperative pain begins when the effects of the local anesthesia wear off and peaks during the first twelve hours after surgery [10]. Third molar surgery is known to cause post-surgical edema or swelling. The swelling typically peaks by the end of the second postoperative day and resolves by the fifth to seventh day. Postoperative events (pain, trismus, swelling) are usually treated with pharmacological drugs and/or other accessory methods such as use of drains, ice packs and laser therapy.

Facial swelling was evaluated by measurements made between 6 known standard reference points: a) Tragus, b) corner of the mouth, c) chin,

d) angle of mandible, e) external canthus and f) bridge of nasal bone. The horizontal measurements are the distances between the outer corner of the mouth and the tragus, as well as the distance from the tragus to the chin region. The distance between the outer canthus of the eye and the angle of the mandible, as well as the tip of the nasal bone to the angle of the mandible, is measured vertically. The facial measurement is determined by the arithmetic mean of the measurements. The pattern of post-surgical pain experienced by the patients in our study was consistent with previous reports; the highest levels of pain were felt on the day of surgery, with a peak 3–5 hours after the surgery [11]. The pain scores were also found to be non-significant.

Proteolytic enzymes are enzymes that are naturally produced by the human body and other living organisms to aid in the performance of essential functions. When these enzymes are supplemented, they have been shown to have an anti-inflammatory effect on body tissues. A study which compared the enzyme complex with a placebo for edema control in orthognathic surgery concluded that usage of the enzymes statistically decreased post-op edema precluding long term corticosteroid use [5].

Chopra et al [12], in their study showed that when compared to betamethasone and ibuprofen, serratiopeptidase had no significant analgesic or anti-inflammatory effect, but it was better than placebo. Al-Khateeb and Nusair et al [13], compared serrapeptase and placebo in reduction of post-op edema and stated that serrapeptase significantly decreases swelling and pain in all subjects, but there was no significant difference in mean maximal interincisal distance between the serrapeptase and placebo groups. Tamimi et al., have also reported recently that serratiopeptidase resulted in a better improvement in the inflammation compared to placebo [14].

Another study which compared bromelain with placebo, found that there were no statistically significant differences between the groups but they noticed a trend towards lesser inflammation and improved mouth opening in the bromelain group [15]. In 2014, Majid and Al-Mashhadani compared bromelain to diclofenac and found that

bromelain improves quality of life following surgical removal of third molars. The investigators used 250 mg of bromelain four times a day in their study [16].

In the current study comparing the enzyme complex and serratiopeptidase, we found that the size of the swelling was almost equal in both groups and was not very significant. The results revealed that there was no significant difference in mouth opening reduction (trismus) between the study groups. This suggests that the enzyme complex is unlikely to be a powerful anti-inflammatory agent. Co-administration of dexamethasone and diclofenac potassium has also been shown to have no effect in reducing postoperative trismus [17].

The combination of both serine (trypsin) and cysteine (Bromelain) proteases is significant as they have different substrate specificities. The mechanism of action of enzymes is not fully understood, but a variety of effects, such as anti-edematous effects and effects on anti-proteinases and α 2 macroglobulin, are thought to contribute to their clinical efficacy [18-23]. The release of inflammatory mediators, the regulation of adhesion molecules, the breaking up of detritus, and the activation of fibrolysis all contribute to better healing with systemic enzyme therapy. Enzymes also decrease immune complexes, which are involved in the pathophysiology of inflammatory rheumatic illnesses [24-29].

CONCLUSION

Within the limitations of our study, it can be concluded that there was no significant difference in the effect of the enzyme combination of rutoside, trypsin and bromelain when compared to serratiopeptidase for reduction of post-operative oedema, pain and trismus following lower third molar surgery. Future studies should be done with higher dosage of the enzymes on a larger sample size to assess the efficacy of these drugs.

Declaration of Interest

The authors report no conflict of interest.

REFERENCES

1. Srivastava N, Shetty A, Goswami RD, Apparaju V, Bagga V, Kale S. Incidence of distal caries in mandibular second molars due to impacted third molars: Nonintervention strategy of asymptomatic third molars causes harm? A retrospective study. *International Journal of Applied and Basic Medical Research*. 2017;7(1):15-19.
2. Halpern LR, Dodson TB. Does prophylactic administration of systemic antibiotics prevent postoperative inflammatory complications after third molar surgery?. *Journal of Oral and Maxillofacial Surgery*. 2007 Feb 1;65(2):177-85.
3. Sortino F, Cicciù M. Strategies used to inhibit postoperative swelling following removal of impacted lower third molar. *Dental research journal*. 2011 Oct;8(4):162.
4. Markovic A, Todorovic LJ. Effectiveness of dexamethasone and low-power laser in minimizing oedema after third molar surgery: a clinical trial. *International journal of oral and maxillofacial surgery*. 2007 Mar 1;36(3):226-9.
5. Shetty V, Mohan A. A prospective, randomized, double-blind, placebo-controlled clinical trial comparing the efficacy of systemic enzyme therapy for edema control in orthognathic surgery using ultrasound scan to measure facial swelling. *Journal of Oral and Maxillofacial Surgery*. 2013 Jul 1;71(7):1261-7.
6. Chandanwale A, Langade D, Sonawane D, Gavai P. A randomized, clinical trial to evaluate efficacy and tolerability of trypsin: chymotrypsin as compared to serratiopeptidase and trypsin: bromelain: rutoside in wound management. *Advances in therapy*. 2017 Jan 1;34(1):180-98.
7. Saghaei M. Random allocation software for parallel group randomized trials. *BMC medical research methodology*. 2004 Dec;4(1):1-6.
8. Janjua OS, Baig Z, Manzoor A, Abbas T. Accuracy of Pederson and modified Parant scale for predicting difficulty level of mandibular third molars. *Arch Orofac Sci*. 2013 Jan 1;8(1):9-13.
9. Baqain ZH, Karaky AA, Sawair F, Khaisat A, Duaibis R, Rajab LD. Frequency estimates and risk factors for postoperative morbidity after third molar removal: a prospective cohort study. *Journal of oral and Maxillofacial Surgery*. 2008 Nov 1;66(11):2276-83.
10. Seymour RA, Blair GS, Wyatt FA. Post-operative dental pain and analgesic efficacy. Part I. *Br J Oral Surg*1983;21: 290–297.
11. Fisher SE, Frame JW, Rout PG, McEntegart DJ. Factors affecting the onset and severity of pain following the surgical removal of unilateral impacted mandibular third molar teeth. *Br Dent J* 1988; 164: 351–354.
12. Chopra D, Rehan HS, Mehra P, & Kakkar AK. A randomized, double-blind, placebo-controlled study comparing the efficacy and safety of paracetamol, serratiopeptidase, ibuprofen and betamethasone using the dental impaction pain model. *Int J Oral Maxillofac Surg* 2009;38:350–355.
13. Al-Khateeb TH, Nusair Y. Effect of the proteolytic enzyme serrapeptase on swelling, pain and trismus after surgical extraction of mandibular third molars. *Int J Oral Maxillofac Surg* 2008;37:264–268.
14. Tamimi Z, Al Habashneh R, Hamad I, Al-Ghazawi M, Kharashgeh H. Efficacy of serratiopeptidase after impacted third molar surgery: a randomized controlled clinical trial. *BMC oral health*. 2021 Dec;21(1):1-9.
15. de la Barrera-Nunez M, Yanez-Vico R, Batista-Cruzado A, Heurtebise-Saavedra J, Castillo-de Oyague R, Torres-Lagares D. Prospective double-blind clinical trial evaluating the effectiveness of Bromelain in the third molar extraction postoperative period. *Med Oral Patol Oral Cir Bucal*. 2014 ;19:e157-162.
16. Majid OW, Al-Mashhadani BA. Perioperative bromelain reduces pain and swelling and improves quality of life measures after mandibular third molar surgery: a randomized, double-blind, placebo-controlled clinical trial. *J Oral Maxillofac Surg*. 2014 ;72:1043-1048.
17. Bamgbose BO, Akinwande JA, Adeyemo WL, Ladeinde AL, Arotiba GT, Ogunlewe MO. Effects of co-administered dexamethasone and diclofenac potassium on pain, swelling and trismus following third molar surgery. *Head Face Med* 2005 7;1:11.
18. Shah D, Mital K. The Role of Trypsin:Chymotrypsin in Tissue Repair. *Adv Ther*. 2018;35(1):31-42.
19. Klein G, Kullich W, Schnitker J, Schwann H. Efficacy and tolerance of an oral enzyme combination in painful osteoarthritis of the hip. A double-blind, randomised study comparing oral enzymes with non-steroidal anti-inflammatory drugs. *Clin Exp Rheumatol* 2006 ;24:25-30.
20. MP SK. Knowledge, Attitude and practices regarding needlestick injuries among dental students. *Asian J Pharm Clin Res*. 2016;9(4):312-5.
21. SK M. Knowledge, attitude, and practices regarding infection control among undergraduate dental students. *Asian J Pharm Clin Res*. 2016;9(1):220-4.

22. Ak H. Knowledge and awareness about oral cancer among undergraduate dental students. Asian Journal of Pharmaceutical and Clinical Research. 2016 Jul 1;165-7.
23. Gayathri MM. Knowledge and awareness among patients about dental implants. Journal of Pharmaceutical Sciences and Research. 2016 May 1;8(5):351.
24. Vijayalakshmi B, Kumar MS. Knowledge of students about Local anaesthetics used during oral surgical procedures. Journal of Pharmaceutical Sciences and Research. 2015 Nov 1;7(11):1011.
25. Gayathri MM. Knowledge, Awareness and Attitude among dental students about hepatitis B infection. Journal of Pharmaceutical Sciences and Research. 2016 Mar 1;8(3):168.
26. Ahamed A, Kumar MS. Knowledge, attitude and perceived confidence in handling medical emergencies among dental students. Journal of Pharmaceutical Sciences and Research. 2016 Jul 1;8(7):645.
27. Kumar S. Knowledge, attitude and practices of dental students toward dental management of patients on antiplatelet therapy. Asian J Pharm Clin Res. 2016;9(30):270-6.
28. MP SK. Local hemostatic agents in the management of bleeding in oral surgery. Asian J Pharm Clin Res. 2016;9(3):35-41.
29. Kumar MP. Newer methods of extraction of teeth. Int J Pharm Bio Sci. 2015;6(3):679-85.

TABLE 1: Mean of Four Distances (mm) Measured Over the Swelling Using Tape Measure Method among the treatment groups.

Treatment groups	Day 0	Day 1	Day 3	Day 7
Serratiopeptidase n=31	52.9 ± 3.4	54.7 ± 3.4*	55.1 ± 3.0*	52.9 ± 3.4 # ^
Enzyme combination n=32	51.5 ± 3.2	53.2 ± 3.2*	54.2 ± 3.3* #	51.9 ± 3.3* # ^
p value	0.40	0.34	0.17	0.20

Data expressed as mean ± SD, p value within group: * p<0.001 compared to Day 1; # p<0.001 compared to Day 3; ^ p<0.001 compared to Day 7 [Student's 't' test]

TABLE 2: The Mean Pain Score Measured Using Visual Analog Scale among the treatment groups.

Treatment groups	Day 0	Day 1	Day 3	Day 7
Serratiopeptidase n=31	8.3 ± 0.9	7.2 ± 0.8*	5.5 ± 1.6* #	2.4 ± 1.7* # ^
Enzyme combination n=32	8.2 ± 0.5	7.3 ± 0.9*	5.4 ± 1.2* #	2.4 ± 1.1* # ^
p value	0.69	0.83	0.88	0.99

Data expressed as mean ± SD, p value within group: * p<0.001 compared to Day 1; # p<0.001 compared to Day 3; ^ p<0.001 compared to Day 7 [Student's 't' test]

TABLE 3: Mean of Mouth Opening (mm) Measured Using Calipers among the treatment groups.

Treatment groups	Day 0	Day 1	Day 3	Day 7
Serratiopeptidase n=31	44.4 ± 3.7	23.1 ± 6.7	29 ± 7.3	35.8 ± 8.2
Enzyme combination n=32	43.4 ± 8.2	21 ± 5.7	24.4 ± 8.2	31.3 ± 9.4
p value	0.62	0.29	0.08	0.12

Data expressed as mean ± SD [Student's 't' test]




FIGURE 1: Requirement for Rescue Medications among the treatment groups.