



## Serum Selenium Levels in Type 2 Diabetes Mellitus Patients

Baneen Sami Tarif<sup>1</sup>, Zainab Hussein AL-Hellawi<sup>2\*</sup>

<sup>1,2</sup>Department of Chemistry, Faculty of Science, Kufa University, Najaf, Iraq

\*Corresponding author: Zainab Hussein AL-Hellawi, Department of Chemistry, Faculty of Science, Kufa University, Najaf, Iraq, Email: Zainab.alhillawi@uokufa.edu.iq

Submitted: 22 February 2023; Accepted: 18 March 2023; Published: 04 April 2023

### ABSTRACT

Diabetes mellitus type 2 (T2DM) is a long-term metabolic disorder characterized by high blood glucose levels. T2DM is caused by insufficient insulin secretion by pancreatic  $\beta$ -cells and an inability of insulin-sensitive tissues to respond appropriately to insulin. In 2019, there are over 463 million people worldwide who have T2DM. Sixty T2DM patients and thirty healthy people served as controls. In terms of age and body mass index (BMI), these patients were comparable to a control group. This study determined demographics and serum selenium (Se). The findings of this study indicated a significant decrease in serum Se in T2DM patients compared to the healthy group. Many T2DM complications are caused by uncontrolled blood glucose and a decrease in antioxidant agents. Increased HbA1c is associated with an increased risk of serious microvascular and macrovascular complications. Low Se levels, on the other hand, increase reactive oxygen species (ROS), and Se deficiency is linked to a number of cardiovascular diseases, including myocardial infarction, cardiomyopathy, atherosclerosis, and coronary heart disease.

**Keywords:** *Selenium, oxidative stress, hyperglycemia and cardiovascular diseases*

### INTRODUCTION

Type 2 Diabetes Mellitus (T2DM) is a broad term for a variety of metabolic disorders characterized by chronic hyperglycemia. The causes are insufficient insulin secretion by pancreatic-cells and an inability of insulin-sensitive tissues to respond appropriately to insulin (Galicja-Garcia et al 2020). By 2019, the number had risen to over 463 million people, and by 2030, it is expected to increase 578 million (Saeedi et al 2019). Diabetes patients usually metabolic disorders of nutrients like glucose, fat, and protein due to hyperglycemia and insulin resistance. In  $\beta$ -cells, glucose metabolism is necessary for the production of insulin, and highly activated glucose metabolism results in the production of reactive

oxygen species (ROS), which reduces insulin secretion. The dual effects of ROS may be related to this contradictory mechanism (Mukai et al 2022). Contrarily, it has been discovered that diabetes is linked to lower levels of intracellular antioxidants like glutathione or vitamins C and E (Rains et al 2011). In several studies, it was discovered that the levels of oxidative stress (OS) biomarkers linked to T2DM had increased significantly and abnormally (Banik et al 2021) induced overproduction of ROS and OS contributes significantly to endothelial dysfunction and is associated with the pathogenesis and development of metabolic diseases (Jasvinder et al 2022). Endothelium is involved in nutrient and metabolite transport and forms an interface between blood

and adjacent tissues in cardiovascular physiology (Alexander et al 2021). When compared to the general population, adults with T2DM are thought to have a 2- to 4-fold higher risk of developing cardiovascular disease (CVD) and passing away (Harding et al 2019). Several metabolic processes, including defense against oxidative stress and healthy cardiovascular function, require selenium (Se), an important trace element. Se plays an essential function in cardiovascular health, as is generally acknowledged (Shimada et al 2021). Se combats free radicals and lowers OS, both of which are thought to improve insulin resistance (Eguchi, et al 2021). Se are essential for the health of the human endocrine, cardiovascular, nervous, and muscle systems as well as the immune and central nervous systems (Roman et al 2014). Prolonged Se deficiency in humans causes serious illnesses because it negatively impacts the functioning of the cardiovascular system and can be a direct cause of myocardial infarction (Shahid et al 2018).

## MATERIALS AND METHODS

### *Subjects*

#### *Patients*

33 women and 27 men who had T2DM were selected. They are between the ages of 50 and 65, beginning in September 2022 and ending in January 2023. All samples were gathered at the Al-Najaf Hospital in the Iraqi province of Najaf. With fasting plasma glucose (FBG) levels of 7.0 mmol/L and HbA1c levels greater than 6.5%, all patients were diagnosed with diabetes according to WHO guidelines. Based on their medical histories, patients were assessed to determine if they had any systemic disorders that might have an impact on the parameters being studied. Patients with thyroid disease, kidney disease,

liver disease, heart disease, or elevated CRP were disqualified from the study.

### *Control group*

Thirty subjects were formed—13 men and 17 women—all of whom appeared to be in good health. Their ages matched those of the patients under investigation. Healthy groups with negative or elevated CRP levels were excluded from the study.

### *Anthropometric values*

Body mass index (BMI), which was calculated using the following formula:  $BMI (kg/m^2) = \text{Weight (kg)} / (\text{Height})^2 (m^2)$ , was used to define obesity. Adults will be categorized in this study as being underweight (BMI 18.5 kg/m<sup>2</sup>), normal weight (BMI 18.5-24.9 kg/m<sup>2</sup>), overweight (BMI 25.0-30.0 kg/m<sup>2</sup>), or obese (BMI > 30.0 kg/m<sup>2</sup>).

### *Biochemical Investigations*

Blood specimens A 5 mL sample of venous blood from each patient or control was drawn using plastic syringes with single-use needles. Two tubes were used to separate the samples. 2 ml of recently drawn venous blood were collected in an EDTA tube to determine the HbA1c level. In order to separate the serum, 3 ml of blood was centrifuged at 3000 rpm for ten minutes after being left at room temperature for 30 minutes to allow for clotting. The serum was then transferred to a fresh, disposable plain tube (hemolysis samples were discarded) and kept in the freezer (-20 °C) until analysis

### *Experimental Apparatus*

**TABLE 1:** Laboratory apparatus and equipment

Instruments	Company	Origin
Centrifuge	Hettich	Germany
Deep freezer	Hitachi	Japan
ELIA Microplate reader	BioTek	USA
Glass gel tube	Q.L.lab	China
Micropipette	Dragon	China

Refrigerator	Hitachi	Japan
Spectrophotometer	Spetra721	Taiwan

### Chemicals

**TABLE 2:** Chemicals and kits used in the study.

Type of Kits	Company/Country
Blood Glucose Kit	Spinreact/Spain
CRP	LTD/Britain
HbA1c	BT LAB/ China
Selenium Kit	BT LAB/ China

### Statistical Analysis

The Kolmogorov-Smirnov test was employed to look into the results group's distribution types. Variable results are classified as either nonparametric or normally distributed according to a statistical distribution. The results were presented as (mean standard deviation) for the normally distributed variable. The Pooled t-test was used to compare the control, patient, and divided groups in the measured parameters. The

difference between groups is regarded as statistically significant when  $p < 0.05$ . SPSS Statistics Base 26 and IBM-USA were used for all statistical analysis. The data was examined using Microsoft Excel 2016 software.

### RESULTS AND DISCUSSION

Table 3 shows the demographic and clinical data for T2DM and the control group.

**TABLE 3:** displays demographic and clinical information about study participants.

Variable	Patients (60) Mean±SD	Controls (30) Mean±SD	p-value
Age (year)	58.1±8.93	58.13±5.23	0.991
No. of male/female	27/33	13/17	0.882
Duration disease (year)	11.98±7.04	-	-
Smoking (Yes/No)	9/51	8/22	0.222
BMI (kg/m <sup>2</sup> )	28.89±4.49	28.01±4.43	0.6074
FBG (mg/dl)	289.23±80.01	98.33±5.6	<0.0001
HbA1c %	9.02±1.53	5.54±0.59	<0.0001
Family History of diabetes (Yes/No)	32/28	0/30	-
CRP P/N	0/60	0/30	-

**BMI:** Body mass index, **FBG:** Fasting blood glucose, **HbA1c:** Glycated hemoglobin, **p:** positive, **N:** negative

Age and BMI do not significantly differ between patients and controls. Age-related T2DM is on the rise, and being more obese is a major risk factor. In young adults, a higher BMI is also independently linked to a higher risk of developing diabetes. (Chen et al 2018). In T2DM compared to the healthy group, there is an expected significant increase in FBS ( $p < 0.001$ ) and HbA1c. HbA1c testing is regarded as a crucial diagnostic tool for tracking dietary

restrictions and treatment plans during the course of diabetes treatment. (Rodas et al 2022). The HbA1c assay accurately and precisely determines chronic glycemia and is associated with a higher risk of developing diabetes complications. (Nusca et al 2019). When the patients' HbA1c is less than 7%, glycemic control has been attained. (Nuha et al 2023). In this study, the patients had uncontrolled blood sugar, with a mean HbA1c of 8.52±1.53%. The risk of serious microvascular and

macrovascular complications increases with HbA1c. (Imai et al 2021). One of the clinical factors that affects patients' glycemic control is long-term T2DM disease. (Traoré et al 2021). Long-term diabetes is linked to worsening beta cell function, which makes blood glucose control challenging. (Kim et al 2014). When a patient had diabetes for more than seven years, the rate of uncontrolled blood glucose was 1.79 times higher than when the patient had diabetes for less than seven years that the longer a person has diabetes, the higher the rate of uncontrolled blood glucose (Ahn J et al 2021).

Heart-related autonomic neuropathy is more common in people with type 2 diabetes when glycemic control is poor and the disease is chronic. (Nagalakshmi et al 2016). Increased levels of circulating free fatty acids and adipokines brought on by obesity decrease insulin sensitivity, increase reactive oxygen species, and possibly impair insulin secretion. (Vrachnis et al 2012).


Additionally, patients with young-onset obesity may have been exposed to maternal undernutrition or overnutrition early in life, which is linked to an increased risk of

complications from type 2 diabetes (Reinehr et al 2016).

Hyperinsulinemia, hyperglycemia, and inflammation that causes OS all cause vascular remodeling. (Ohishi et al 2018). T2DM patients are more likely to die from any cause and develop cardiovascular disease (CVD), especially if they also have conventional CVD risk factors like hypertension, hyperglycemia, abdominal obesity, and dyslipidaemia. (Rawshani et al . 2018). An Egyptian study on diabetic patients discovered that the patients' level of education, line of work, and smoking habits all affected how well their diabetes was controlled. Also mentioned were factors that affected glycemic control, but these were specific to primary care physicians and included things like living in a rural area, taking part in diabetes training, being older, having dealt with diabetic patients for a longer period of time, and how well they were following the recommendations (Azzam et al 2021).

**Serum Selenium**

Patients with T2DM have significantly lower serum Se levels than the healthy group (p=0.027).



**FIGURE 1:** Serum Se in T2DM patients and Controls in bars plot.

Selenium (Se) is a crucial element for a variety of biological processes, including the protection against OS and inflammation and function as

vital enzymes for redox homeostasis. (Zhang et al 2017). A low level of Se impairs important metabolic processes by increasing ROS and

decreasing mitochondrial function. (Shimada et al 2021). The relationship between Se status and glucose regulation extends to hypoglycemia as well as hyperglycemia. (Wang et al 2020). Numerous studies have suggested altered Se concentrations and increased metabolic needs for Se in patients with diabetes, which is characterized by a hyperglycemic state that may lead to OS and inflammatory activation. (Wang et al 2017). Among adults with T2D, a recent study discovered a linearly inverse relationship between serum selenium concentrations and heart disease mortality. (Zixin et al 2022). Cardiomyopathy, atherosclerosis, and coronary heart disease, as well as heart failure, myocardial infarction, and heart failure, are all linked to se deficiency. (Rayman et al 2008). Se deficiency has been linked to heart disease, including Keshan disease, in other studies (Liu et al 2019). On the other hand, a study found that supplementing with Se may help to prevent heart diseases. (Wang et al 2020). In an animal study, it was discovered that Se therapy may improve mice with atherosclerosis. (Guo et al 2020). Given that T2DM is linked to OS, it was initially anticipated that Se would help diabetic patients. At high supranutritional doses, Se (as selenate) does, in fact, have anti-diabetic and insulin-mimetic effects (Steinbrenner et al 2013). Se performs a variety of tasks in the body's cells, including fighting free radicals and lowering OS, which is thought to improve insulin resistance. (Eguchi et al 2021). Numerous observational studies conducted in various nations found that gestational DM pregnant women had lower Se levels than euglycemic pregnant women. (Moshfeghy et al . 2020).

#### ACKNOWLEDGEMENT

The authors express their thanks to the medical team at "AL-Najaf Teaching Hospital" in Najaf, Iraq, as well as the patients for their cooperation

#### REFERENCES

- Ahn J, Yang Y. Factors Associated with Poor Glycemic Control Amongst Rural Residents with Diabetes in Korea. *Healthcare (Basel)*. 2021 Apr 1;9(4):391. doi: 10.3390/healthcare9040391. PMID: 33915834; PMCID: PMC8065919.
- Alexander Y, Osto E, Schmidt-Trucksäss A, Shechter M, Trifunovic D, Duncker DJ, et al. Endothelial function in cardiovascular medicine: a consensus paper of the European Society of Cardiology Working Groups on Atherosclerosis and Vascular Biology, Aorta and Peripheral Vascular Diseases, Coronary Pathophysiology and Microcirculation, and Thrombosis. *Cardiovasc Res*. (2021) 117:29–42.
- Azzam MM, Ibrahim AA, Abd El-Ghany MI. Factors affecting glycemic control among Egyptian people with diabetes attending primary health care facilities in Mansoura District. *Egypt J Crit Care*. 2021;33:33.
- Banik S, Ghosh A. The association of oxidative stress biomarkers with type 2 diabetes mellitus: A systematic review and meta-analysis. *Health Sci Rep*. 2021 Oct 1;4(4):e389.
- Chen Y, Zhang XP, Yuan J, Cai B, Wang XL, Wu XL, Zhang YH, Zhang XY, Yin T, Zhu XH, Gu YJ, Cui SW, Lu ZQ, Li XY. Association of body mass index and age with incident diabetes in Chinese adults: a population-based cohort study. *BMJ Open*. 2018 Sep 28;8(9):e021768.
- Eguchi, N.; Vaziri, N.D.; Dafoe, D.C.; Ichii, H. The Role of Oxidative Stress in Pancreatic  $\beta$  Cell Dysfunction in Diabetes. *Int. J. Mol. Sci*. 2021, 22, 1509.
- Galicía-García U, Benito-Vicente A, Jebari S, Larrea-Sebal A, Siddiqi H, Uribe KB, Ostolaza H, Martín C. Pathophysiology of Type 2 Diabetes Mellitus. *Int J Mol Sci*. 2020 Aug 30;21(17):6275.
- Guo, L.; Xiao, J.; Liu, H.; Liu, H. Selenium nanoparticles alleviate hyperlipidemia and vascular injury in ApoE-deficient mice by regulating cholesterol metabolism and reducing oxidative stress. *Metallomics* 2020, 12, 204–217
- Harding JL, Pavkov ME, Magliano DJ, Shaw JE, Gregg EW. Global trends in diabetes complications: a review of current evidence. *Diabetologia*. 2019;62(1):3–16.
- Imai C, Li L, Hardie RA, Georgiou A. Adherence to guideline-recommended HbA1c testing frequency and better outcomes in patients with type 2 diabetes: a 5-year retrospective cohort study in Australian general practice. *BMJ Qual Saf*. 2021 Sep;30(9):706-714.
- Jasvinder Singh Bhatti, Abhishek Sehrawat, Jayapriya Mishra, Inderpal Singh Sidhu, Umashanker Navik, Naina Khullar, Shashank Kumar, Gurjit Kaur Bhatti, P. Hemachandra Reddy. Oxidative stress in the pathophysiology of type 2 diabetes and related complications: Current therapeutic strategies and future perspectives., 1 May 2022, Pages 114-134.



12. Kim S.Y. It's still not too late to make a change: Current status of glycemic control in Korea. *J. Diabetes Metab.* 2014;38:194–196. doi: 10.4093/dmj.2014.38.3.194.
13. Liu, X.; He, S.; Tan,W. Expression profile analysis of selenium-related genes in peripheral blood mononuclear cells of patients with Keshan disease. *Biomed Res. Int.* 2019, 2019, 4352905.
14. Moshfeghy, Z.; Bashiri, K.; Dabbaghmanesh, M.H.; Akbarzadeh, M.; Asadi, N.; Sayadi, M. The Predictive Value of Selenium in Diagnosis of Gestational Diabetes: A Nested Case-Control Study. *Int. J. Gen. Med.* 2020, 13, 53–60.
15. Mukai E, Fujimoto S, Inagaki N. Role of Reactive Oxygen Species in Glucose Metabolism Disorder in Diabetic Pancreatic  $\beta$ -Cells. *Biomolecules.* 2022; 12(9):122.
16. Nagalakshmi, V., Smruthi, V., Shama, Rathnakarashetty, G., Nallulwar, S.C., 2016. Cardiac Autonomic Neuropathy in Type II Diabetes Mellitus Patients and its Association with the Age of the Patient, Duration of Disease and Glycemic Control. *Indian journal of physiology and pharmacology* 60, 222-229.
17. Nuha A. ElSayed, Grazia Aleppo, Vanita R. Aroda, Raveendhara R. Bannuru, Florence M. Brown, Dennis Bruemmer, Billy S. Collins, Marisa E. Hilliard, Diana Isaacs, Eric L. Johnson, Scott Kahan, Kamlesh Khunti, Jose Leon, Sarah K. Lyons, Mary Lou Perry, Priya Prahalad, Richard E. Pratley, Jane Jeffrie Seley, Robert C. Stanton, Robert A. Gabbay; on behalf of the American Diabetes Association, 6. Glycemic Targets: Standards of Care in Diabetes—2023. *Diabetes Care* 1 January 2023.
18. Nusca, Annunziata, Dario Tuccinardi, Claudio Proscia, Rosetta Melfi, Silvia Manfrini Antonio Nicolucci, Antonio Ceriello, Paolo Pozzilli, Gian Paolo Ussia, and Francesco Grigioni.. “Incremental Role of Glycaemic Variability over HbA1c in Identifying Type 2 Diabetic Patients with High Platelet Reactivity Undergoing Percutaneous Coronary Intervention.” *Cardiovascular Diabetology* 2019.18 (1): 147.
19. Ohishi, M. Hypertension with diabetes mellitus: Physiology and pathology. *Hypertens. Res.* 2018, 41, 389–393.
20. Rains J.L., Jain S.K. Oxidative stress, insulin signaling, and diabetes. *Free Radic. Biol. Med.* 2011;50(5):567–575.
21. Rawshani A, Rawshani A, Franzen S, Sattar N, Eliasson B, Svensson AM, et al. Risk factors, mortality, and cardiovascular outcomes in patients with type 2 diabetes. *N Engl J Med.* 2018;379(7):633–644.
22. Rayman, M.P. Food-chain selenium and human health: Emphasis on intake. *Br. J. Nutr.* 2008, 100, 254–268 Rayman, M.P. Food-chain selenium and human .
23. Reinehr T, Karges B, Meissner T, et al. Inflammatory Markers in obese adolescents with type 2 diabetes and their relationship to hepatokines and adipokines. *J Pediatr* 2016;173:131–5. doi:10.1016/j.jpeds.2016.02.055
24. Rodas Getachew Abera, Eyouel Shimeles Demesse and Wako Dedecha Boko Evaluation of glycemic control and related factors among outpatients with type 2 diabetes at Tikur Anbessa Specialized Hospital, Addis Ababa, Ethiopia: a cross-sectional study Abera et al. *BMC Endocrine Disorders* (2022) 22:54.
25. Roman M., Jitaru P., Barbante C. Selenium biochemistry and its role for human health. *Metallomics.* 2014;6:25–54.
26. Saeedi P, Petersohn I, Salpea P, Malanda B, Karuranga S, Unwin N, Colagiuri S, Guariguata L, Motala AA, Ogurtsova K, Shaw JE, Bright D, Williams R; IDF Diabetes Atlas Committee. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9th edition. *Diabetes Res Clin Pract.* 2019 Nov;157:107843.
27. Shahid, M.; Niazi, N.K.; Khalid, S.; Murtaza, B.; Bibi, I.; Rashid, M.I. A critical review of selenium biogeochemical behavior in soil-plant system with an inference to human health. *Environ. Pollut.* 2018, 234, 915–934.
28. Shimada BK, Alfulaij N, Seale LA. The Impact of Selenium Deficiency on Cardiovascular Function. *International Journal of Molecular Sciences.* 2021; 22(19):10713.
29. Steinbrenner H. Interference of selenium and selenoproteins with the insulin-regulated carbohydrate and lipid metabolism. *Free Radic. Biol. Med.* 2013;65:1538–1547.
30. Traoré S, Guira O, Zoungrana L, et al. Factors associated with prolonged poor glycemic control in type 2 diabetes mellitus (T2DM) patients followed in the Department of Internal Medicine at the Yalgado Ouedraogo teaching hospital, Ouagadougou (Burkina Faso) *Open J Intern Med.* 2021;11:1–26.
31. Vrachnis N, Antonakopoulos N, Iliodromiti Z, et al. Impact of maternal diabetes on epigenetic modifications leading to diseases in the

- offspring. *Exp Diabetes Res* 2012;2012:1–6. 10.1155/2012/538474).
32. Wang N Tan HY, Li S, Xu Y, Guo W, Feng Y. Supplementation of micronutrient selenium in metabolic diseases: its role as an antioxidant. *Oxid Med Cell Longev.* 2017;2017:1.
33. Wang Y., Rijntjes E., Wu Q., et al. Selenium deficiency is linearly associated with hypoglycemia in healthy adults. *Redox Biol.* 2020;37:101709.
34. Wang, L.; Yin, J.; Guo, X. Serious selenium deficiency in the serum of patients with kashin-beck disease and the effect of nano-selenium on their chondrocytes. *Biol. Trace Elem. Res.* 2020, 194, 96–104.
35. Zhang H, Yan C, Yang Z, Zhang W, Niu Y, Li X, Qin L and Su Q. Alterations of serum trace elements in patients with type 2 diabetes. *J Trace Elem Med Biol* 2017; 40: 91-96.
36. Zixin Qiu, Tingting Geng, Zhenzhen Wan, Qi Lu, Jingyu Guo, Liegang Liu, An Pan, Gang Liu, Serum selenium concentrations and risk of all-cause and heart disease mortality among individuals with type 2 diabetes, *The American Journal of Clinical Nutrition*, Volume 115, Issue 1, January 2022, Pages 53–60.