



## PRELIMINARY PHYTOCHEMICAL ANALYSIS AND *IN VITRO* BIOLOGICAL ACTIVITIES OF *OTOSTEGIA LIMBATA* LEAVES ETHANOLIC EXTRACT AGAINST ORAL PATHOGENS

Khursheed Ur Rahman<sup>1\*</sup>, Ghulam Mujtaba Shah<sup>1</sup>, Muhammad Ajmal Shah<sup>2</sup>, Muhammad Ikram<sup>3</sup>, Muhammad Fiaz<sup>1</sup>, Jan Alam<sup>1</sup>, Muhammad Sajid<sup>4</sup>

<sup>1\*</sup> Department of Botany, Hazara University, Mansehra 21300, Pakistan.

<sup>2</sup> Department of Pharmacy, Hazara University, Mansehra 21300, Pakistan.

<sup>3</sup> Department of Pharmacy, COMSATS University Islamabad, Abbottabad Campus 22060, Pakistan

<sup>4</sup> Department of Agriculture, Hazara University, Mansehra 21300, Pakistan.

\*Corresponding author: Khursheed Ur Rahman

\*Email: khursheed823@yahoo.com

### Abstract

In the present investigation, we have accessed Phytochemicals, HPLC, Antibacterial, Antifungal and Cytotoxic activity of *Otostegia limbata*. It diverted our attention to one of its potent species to be unveiled in this research. The focus of this study was to examine systematically its biological activities and seek out its chemical constituents. Qualitative phytochemical analysis removed phenols, glycosides, flavonoids, alkaloids, quinone, carbohydrates, amino acid, terpenes and coumarins are presence while tannins, saponins and sterols are absence. In Quantitative the *O. limbata* leaves parts for isolation of active phtometabolites namely alkaloids, sterol, flavonoids, tannins and phenols. The HPLC analysis showed that the samples contained eight identified phenolic compounds, of which Gallic acid, Catechol, Hydroxybenzoic acid, Caffeic acid were most abundant. The antibacterial activity of ethanolic extract displayed the highest inhibition region against *Streptococcus mitis* that was  $35 \pm 0.1$  (ZOI $\pm$ SD) and the lowest inhibition region against *S. aureus* that was  $14 \pm 0.3$  (ZOI $\pm$ SD) in 200mg/ml  $\pm$  SD. The antifungal activity resulted that ethanol shows the maximum inhibition zone against *A.fumigatus* that was  $29 \pm 0.1$  (ZOI $\pm$ SD) and the minimum inhibition zone against *Aspergillus flavus* that was  $19 \pm 0.4$  (ZOI $\pm$ SD) in 200mg/ml  $\pm$  SD. The ethanolic extract of research plant were exposed to cytotoxic assay at concentrations is 50 $\mu$ g/ml, 100 $\mu$ g/ml and 150 $\mu$ g/ml and their results were calculated that indicates that at 50 $\mu$ g/ml is 60%, 100 $\mu$ g/ml is 70% and 150 $\mu$ g/ml is 80%. The current study suggested that, after the isolation of individual components, *O. limbata* be investigated for assessing biological activity.

**Key word:** Phytochemicals, HPLC, Antibacterial, Antifungal and Cytotoxic activity, *Otostegia limbata*.

**Key findings:** The selected medicinal plants *Otostegia limbata* which is used in the tooth pain we are chick the effect of antimicrobial activity against oral pathogens, effect of cytotoxic activity against Brime shrimps and also check the photochemical screening tests.

## 1. INTRODUCTION

Pakistan has received a priceless gift from nature in the form of medicinal plants. Since ancient civilizations, medicinal plants have occupied a permanent position for treating a variety of diseases (Anand *et al.*, 2019). Natural materials and their preparations make a significant contribution to solving practical issues for people, animals, agricultural, veterinary, food goods, cosmetics, and other industries (Drasar and Khripach, 2019). The family Lamiaceae includes one of the well-known genera *Otostegia*, which is geographically widespread around the world. There are roughly 4000 species and 220 genera in this group of flowering plants. *Otostegia limbata*, also known as *Rydingia limbata* (Benth.) Scheen & V. A. Albert and *Ballota limbata* is a significant medicinal plant of this genus (Scheen and Albert, 2017). The common names "Spin aghzai," "Chiti booti," "Chitti jharri," "Spin azghay," and "Bui" are used to identify certain plant species.. The plant's distinctive features include a cluster of pale yellow flowers, oblong leaves with a thick pointed tip, pointy bracts, and a tiny petiole. Plant species contain a variety of chemical components, including the acids ballotenic and ballodiolic, limbatolide A, B, C, and D (Sadaf *et al.*, 2016). The *Otostegia limbata* is commonly used as an ethnomedicine in Pakistan for a variety of ailments, including jaundice, cancer, scabies, boils, goitre, ulcer, cuts, wounds, dental issues, and animal diseases (Rosselli *et al.*, 2019). Traditional healers use fresh leaf infusion to treat conditions including acidity, hypertension, depression, ulcer, jaundice, gum disease, and ocular infection in Pakistan's Azad Jammu & Kashmir, KPK, Punjab, and Himalayan regions (Rashid *et al.*, 2015). To meet our everyday basic needs, nature has bestowed upon us a wealth of valuable treasures. Plants are among the most crucial sources. Plants have been used for therapeutic purposes for a very long time. These ancient medicinal plants are essential to complementary medicine since they are used to cure a variety of ailments. The Indian subcontinent is home to a rich trove of various plant species with a variety of practicable medicinal characteristics. Herbal medicines also play a major role for gums and oral problems. Herbal medications have unique recommendations and a long history of respectability. Herbal medicine, which was once used to treat heart conditions like heart failure, plays a vital part in the management and treatment of disorders like digitalis, which contains cardiac glycosides (Arora and Arora, 2021). Even in this cutting-edge, technological age, doctors still recommend a variety of medications with botanical origins. Up to 10% of local communities around the world employ medicinal plants to treat various illnesses, yet only 1% of these plants have been identified by scientists. The Alkaloids, tannins, and flavonoids, among other secondary metabolites, are widely distributed in plants with antibacterial characteristics. Because medicinal plants are less poisonous and have less negative effects, they are utilised to treat a variety of ailments (Morrison *et al.*, 1980). The development of caries and periodontal illnesses is significantly influenced by germs on the tooth surface, according to the movie Dental Plaque (Gamboe *et al.*, 2008). Mutans streptococci have the capacity to produce extracellular polysaccharides from sucrose, mostly water-insoluble glucan, using the glucosyl transferase enzyme, which allows them to colonise the tooth surface and start the production of plaque (Bankova *et al.*, 1992). This sucrose-dependent adherence and accumulation of cariogenic streptococci is important to the establishment of a pathogenic plaque. The microbial composition of the plaque surrounding the gingival margin and subgingival area may change from being dominated by streptococcus to being more Actinomyces species and more capnophilic and necessary anaerobic bacteria, including Porphyromonas gingivalis (Aga *et al.*, 1994). These microbes appear to play a role in periodontal disease and root caries, respectively. Therefore, antimicrobial treatments for certain oral pathogens, especially those that might alter plaque production, could be very effective in preventing dental caries and periodontal disorders. The plant's ethanolic extract has numerous pharmacological properties, including anti-inflammatory, anaesthetic, and cytostatic effects in addition to antibacterial activity. Streptococcus mutans is another bacterium that it is antibacterial for (Koo *et al.*, 2000). There is, however, little information available regarding its antibacterial efficacy against other oral pathogens or its impact on dental plaque formation in vitro. Over the past few decades, there has been a noticeable growth of bacteria that are resistant to antibiotics. Antibiotic overuse and abuse are the primary causes of the rising

prevalence of resistant microorganisms worldwide. It's interesting to note that traditional medicine, including herbal medicine, has long been used in developing nations for healthcare and numerous studies have confirmed its efficacy in controlling a variety of infectious diseases (WHO, 2002). Plant extracts made from the leaves, stems, and roots serve as a valuable resource for the discovery of powerful and innovative antibacterial and biofilm medications (Essawi and Srour, 2000). *Otostegia limbata* is a spiny, 40–60 cm tall shrub with many branches (Fig. 1). It is known as "spin azghay" locally in (Lower Dir) and thrives in dry environments. It is widely grown in Kashmir and throughout Pakistan. *O. limbata* is useful for treating wounds and is effective against ophthalmia, gum, and skin problems (Hedge *et al.*, 1990). This study focuses on the crude methanolic extracts, water, and hexane fractions of aerial portions (leaves) from *J. regia* and *O. limbata*'s anti-pseudomonal activity against *P. aeruginosa* planktonic and biofilm forms *in vitro* (Kale *et al.*, 2011) and (Abbasi *et al.*, 2010).



**Figure 1a.** *Otostegia limbata* (Benth.) Boiss



**Figure 1b.** Herbarium specimen

## 2. MATERIALS AND METHODS

### 2.1 Collection and authentication of plants

The plant sample was collected from district (Mansehra) during session June 2021 and Identified with the help of flora and taxonomist expert by Prof. Dr Ghulam Mujtaba Shah, Chairman, Department of Botany, Hazara University Mansehra KP, Pakistan. After identification the voucher Number (15060) was assigned to the plant species and specimen were deposited in the Herbarium of Hazara University (HUP) for permanent record (Fig. 2). The plant materials were washed with tap water, separated and dried in shade for 15 days. These materials were used afterward for phytochemical, and biological activities of *in-vitro* biological screening i-e antimicrobial activities against oral pathogen and cytotoxic potential activity. The plant material was powdered with the help of electrical grinder. The Whatman filter paper was used after the muslin cloth to filter the extracts. Rotary evaporation will be used at 40°C to remove extra solvent from the filtrate. Until further examination, the extract was kept in a container of amber colour.

## 2.2 Extraction of plants material

Plant extracts were prepared using microwave extraction technology, according to a previously reported procedure. The microwave's power setting was set at 9000 W. There are three basic stages to this process. In the first step, 750 mL of ethanol and 100 g of each plant powder were added to separate beakers in a 1000 mL container. The microwave was on for 2 minutes, then off for 30 seconds while the beakers were in it. Five times these procedures were carried out. The same process will be used to complete two additional cycles. The muslin cloth will be used to filter the extracts first, and then Whatman filter paper. Rotary evaporation at 40°C was used to remove extra solvent from the filtrate. The extracts will be kept till further analysis in a container of amber colour (Farrukh *et al.*, 2022).

## 2.3 Phytochemical Analysis of *Otostegia limbata*

### 2.3.1 Qualitative analysis

Different protocols used to detect the presence or absence of different classes of phytochemicals (Phlobatannins, Cardiac glycosides, Quinones, Steroids, Saponins, Coumarins, Tannins and Terpenoids). Presence of these chemicals was detected by production of different colours (Chung *et al.*, 1998).

### 2.3.2 Quantitative analysis

The spectrometere was used in quantitative phytochemical analyses of total alkaloid contents, total saponins, total flavonoids contents, total tannins contents and total phenolic contents by using standered protocol methods (Haq *et al.*, 2016).

## 2.4 High performance liquid chromatography (HPLC) analyses

In 20 ml of methanol (62.5%) and 5 ml of HCl were used to extract the ground plant material (6M). The extract was sonicated for 15 minutes and then refluxed in a water bath for two hours following nitrogen purging (Muhammad *et al.*, 2020). Before injecting into HPLC, filter the extract twice via a 0.2 m Millex-HV membrane filter. The Shimadzu LC-20AT HPLC system includes a column oven, an auto-sampler, and a diode array detector (SPD-M20A). Utilized was an analytical column with a guard column (KJO-4282, Phenomenex): Purospher Star RP-18 endcapped 5 m 100 Å (250 x 4.60 mm, Merck, Germany). The composition gradient programme was used with just minor alterations with the mobile phase consisting of (A) 0.1% acetic acid and (B) methanol (Qasim *et al.*, 2017). The flow rate was 0.8 mL per minute. By contrasting the retention times and UV-Vis spectra of chromatographic peaks with those of genuine reference standards at 280 nm, phenolic chemicals were identified (Jamshed *et al.*, 2019).

## 2.5 Antimicrobial activity

Anti-microbiological action 100 mg/ml, 150 mg/ml, and 200 mg/ml of crude extract were the chosen concentrations. Standard antibiotics were employed as the drug of choice for the positive control for various bacterial and fungal infections, and DMSO was utilised for the negative control. Drugs that were in powder form had been accurately weighed and dissolved in the proper dilutions to the necessary 200 mg/mL concentration. The antibacterial assays were evaluated using the agar-well diffusion method. Agar Mueller-Hinton was employed to prepare the media (Enjalric *et al.*, 1987).

### Test organisms

The Six bacterial strains (*Streptococcus mutans* (ATCC 25175), *Streptococcus mitis* (ATCC 23175), *Staphylococcus aureus* (ATCC-6538), *Pseudomonas aeruginosa* (ATCC- 15442, *Bacillus subtilis* (ATCC6633), and *Escherichia coli* (ATCC-25922), associated with dental infections were used for antibacterial analyses. Similarly *Aspergillus flavus* (FCBP-0064), *Aspergillus fumigatus* (FCBP-66), *Aspergillus niger* (FCBP-0198), *Fusarium solani* (ATCC 36031) and *Candida albicans* (ATCC

26081) were used to detect antifungal activity. Cephadrine 50µg was used as positive control for antibacterial activity and same quantity of fluconazole used as positive control against fungal strains. DMSO was used negative control against bacterial and fungal strains. The Department of Microbiology at Hazara University in Mansehra, KPK, Pakistan and the Department of Biotechnology at the University of Science and Technology in Bannu provided all the microorganisms. Throughout the study, stock cultures of bacteria and fungi were kept in their proper growth medium at 4 °C. Both antibacterial and antifungal studies were conducted using the agar well diffusion method (Carron *et al.*, 1987).

## 2.6 Cytotoxic activity

The cytotoxic activity was done by following standard protocol method (Meyer *et al.*, 1982).

### Required media

Brine shrimp eggs, sea salt, distilled water, a tray or container with partitions, plant extract, test tubes, micro tips, and a magnifying glass.

### Stock solution preparation

The 20 mg of Plant extract were dissolved in 2 ml of ethanol to create the stock solution.

### Method

Following techniques allowed for the determination of the plant's potential for cytotoxicity. Brine shrimp eggs were first placed in a plastic container or tray with a perforated partition and 3.8 grammes of sea salt was first dissolved in 1000 ml of distilled water. This media was then added, and the container was placed at a temperature of 34–36 oC for one day to hatch the brine shrimp eggs. As they emerged, the shrimp went to the opposite side of the container. Following the creation of three concentrations 100 mg/ml, 500 mg/ml, and 1000 mg/ml stock solutions were added in accordance with these concentrations and the test tubes were left for the remainder of the day to allow the ethanol to evaporate. Next, 2 mg of sea salt was added to the test tubes to make the total volume 5 mg, and ten newly hatched brine shrimp were then placed inside the test tubes using a micro-pipette, and the tubes were then left The following day, using a microscope, the number of alive and dead brine shrimp in each test tube was determined.

### Statistical analyses

Data tabulated and analyzed by using statistic software statistic 8.1.

## 3. RESULTS

### 3.1 Qualitative phytochemical analysis of *Otostegia limbata*

Phytochemical analysis revealed that the crude extract of *Otostegia limbata* included many different types of compounds such as phenolic and glycosides, as well as flavonoids and alkaloids, as well as quinones, carbohydrates, amino acid, terpenoids and coumarins. However, the tannins, saponins and sterols test results for the crude extract showed no change in colour (Table No.1)

**Table No 1. Qualitative phytochemical analyses of ethanolic extract of *Otostegia limbata***

S.No	Constituents	Present (+)	Absent (-)
1	Phenols	+	
2	Glycosides	+	
3	Tannins		-
4	Flavonoids	+	
5	Alkaloids	+	
6	Saponins		-
7	Quinones	+	
8	Sterols		-

9	Carbohydrates	+	
10	Amino acids	+	
11	TeT Terpenoids	+	
12	Coumarins	+	

Key = Negative sign (-) indicate absence, positive sign (+) indicate presence

### 3.2 Quantitative analysis of *O. limbata*

Quantitative phytochemical screening of the leaves parts of the *O. limbata* for isolation of active phtometabolites namely alkaloids, sterol, flavonoids, tannins and phenols in (Table No.2).The results revealed the bioactive constituents in leaves are alkaloids were in the range of ( $16.66 \pm 1.33$  mg/g) and sterol ( $14.68 \pm 0.66$  mg/g), flavonoids ( $11.5 \pm 0.33$  mg/g), tannins is ( $14.30 \pm 0.10$  mg/g) and phenols is ( $56.73 \pm 0.25$  mg/g).

**Table 2. Quantitative analysis of *Otostegia limbata*. All values are mean  $\pm$  SEM of three determinations. All values are expressed in mg/g.**

S.No	Extract	Alkaloids	Sterol	Flavonoids	Tannins	Phenol
1	OLL	$16.66 \pm 1.33$	$14.68 \pm 0.66$	$11.5 \pm 0.33$	$14.30 \pm 0.10$	$56.73 \pm 0.25$

Key = OLL= *Otostegia limbata* Leaves, ND= Not detected


### 3.3 High performance liquid chromatography (HPLC) analyses

The HPLC analysis showed that the samples compound is contained eight identified phenolic compounds, of which Gallic acid, Catechol, Hydroxybenzoic acid, Caffeic acid were most abundant. The Gallic acid is  $12.11 \pm 0.25$ , Catechol is  $3.01 \pm 0.07$  and Caffeic acid  $7.57 \pm 0.92$ . The highest compound is Gallic acid is  $12.11 \pm 0.25$  while the lowest compound is Catechol is  $3.01 \pm 0.07$

**Table 3. Phenolic composition (mg g<sup>-1</sup> dry weight) of *Otostegia limbata***

S.No	Compounds	Retention time	<i>O. limbata</i>
1	Pyrogallol	12.557	n.d
2	Gallic acid	15.192	$12.11 \pm 0.25$
3	Catechol	18.145	$3.01 \pm 0.07$
4	Hydroxybenzoic acid	19.136	n.d
5	Chlorogenic acid	23.59	n.d
6	Caffeic acid	24.756	$7.57 \pm 0.92$
7	Coumaric acid	30.039	n.d
8	Ferulic acid	31.835	n.d





**Figure 2.** HPLC chromatograms of standard compounds *i.e.* Pyrogallol (1), Gallic acid (2), Catechol (3), Hydroxybenzoic acid (4), Chlorogenic acid (5), Caffeic acid (6), Coumaric acid (7), and Ferulic acid (8) and *Otostegia limbata* extract.


### 3.4 Antibacterial activity of *Otostegia limbata*

Figure 2 displays the *Otostegia limbata* ethanolic extract's antibacterial activity. *Streptococcus mitis* showed a maximum inhibition zone of 35 0.1, while *S. aureus* showed a minimum inhibition zone of 14 0.3 in 200 mg/ml standard deviation. Antibiotics had a maximum zone of inhibition of 45 0.4 against *P. aerogenose* and a minimum zone of inhibition of 28 0.5 against *S. aureus* in 100 mg/ml SD. Table 3 of the results shows the results for the plant extract and antibiotic in terms of standard and mean deviation values. When used as a negative control, DMSO does not inhibit bacterial strain development (Table No. 4).

**Table 4. Antibacterial activity of *Otostegia limbata***

Tests Microorganism	Antibiotics ZOI(mm) ± SD	Leaves ZOI(mm) Means ± SD		
		OLLEE 100mg/ml ± SD	OLLEE 150mg/ml ± SD	OLLEE 200mg/ml ± SD
<b>Bacterial strains</b>	<b>Cephradine 100mg/ml ZOI(mm)</b>			
<i>S. mutans</i>	38 ± 0.1	18 ± 0.2	20 ± 0.6	22 ± 0.4
<i>S. mitis</i>	42 ± 0.3	14 ± 0.6	29 ± 0.2	35 ± 0.1
<i>S. aureus</i>	28 ± 0.5	8 ± 0.4	10 ± 0.5	14 ± 0.3
<i>P. aerogenose</i>	45 ± 0.4	25 ± 0.5	29 ± 0.3	32 ± 0.5
<i>B. substilus</i>	43 ± 0.6	19 ± 0.3	20 ± 0.1	22 ± 0.2
<i>E. coli</i>	40 ± 0.2	17 ± 0.1	24 ± 0.4	32 ± 0.6

**Key=** *S. mutans*= *Streptococcus mutans*, *S. mitis*= *Streptococcus mitis*, *S. aureus* = *Staphylococcus aureus*, *P. aerogenose*= *Pseudomonas aeruginosa*, *B. substilus*= *Bacillus subtilis*, *E. coli* = *Escherichia coli* and **OLLEE**= *Otostegia limbata* leaf ethanolic extracts, **SD** =Standard deviations)



Key= PSLEE (*Otostegia limbata* leaves ethanolic extract)

Figure 3. Graphical representation of antibacterial activity of *Otostegia limbata*

### 3.5 Antifungal activity of *Otostegia limbata*

The greatest inhibition zone against *Aspergillus fumigatus* was 29 0.1, and the minimum inhibition zone against *Aspergillus flavus* was 19 0.4, according to an ethanol extract of the antifungal activity. Antibiotics had a maximum zone of inhibition of 36.66 0.4 against *C. albicans* and a minimum zone of inhibition of 29.66 0.2 against *A. niger* in (table 4) of the results shows the results for the plant extract and antibiotic in terms of standard and mean deviation values. DMSO is utilised as a negative control and exhibits no growth suppression or resistance to fungi. (Table 5).

Table 5. Antifungal activity of ethanolic extracts of *Otostegia limbata*

Tests	Antibiotics	Leaves ZOI(mm) Means $\pm$ SD		
Microorganism	ZOI(mm) $\pm$ SD			
Fungal strains	Fluconazole 100mg/ml ZOI(mm)	OLLEE 100mg/ml $\pm$ SD	OLLEE 150mg/ml $\pm$ SD	OLLEE 200mg/ml $\pm$ SD
<i>F. flavus</i>	35.33 $\pm$ 0.5	10 $\pm$ 0.2	12 $\pm$ 0.6	19 $\pm$ 0.4
<i>A. fumigatus</i>	32.33 $\pm$ 0.3	12 $\pm$ 0.6	15 $\pm$ 0.2	29 $\pm$ 0.1
<i>C. albicans</i>	36.66 $\pm$ 0.4	13 $\pm$ 0.4	15 $\pm$ 0.5	19 $\pm$ 0.3
<i>A. niger</i>	29.66 $\pm$ 0.2	16 $\pm$ 0.5	20 $\pm$ 0.3	21 $\pm$ 0.5
<i>F. solani</i>	30.33 $\pm$ 0.1	12 $\pm$ 0.3	15 $\pm$ 0.1	19 $\pm$ 0.2

Key= *F. flavus* = *Aspergillus flavus*, *A. fumigatus* = *Aspergillus fumigatus*, *C. albicans* = *Candida albicans*, *A. niger* = *Aspergillus niger*, *F. solani* = *Fusarium solani* and OLLEE= *Otostegia limbata* Leaf ethanolic extracts, S.D =Standard deviations)






Figure 4. Graphical representation of antifungal activity of *Otostegia limbata*


### 3.6 Cytotoxic Brine shrimps assay

The cytotoxic activity of *O. limbata* extracts at various doses (50, 100 and 150µg/ml) was conducted. It was shown that *O. limbata* extracts had a cytotoxic impact on brine shrimps when evaluated for 72 hours under controlled conditions. Results after 24 hours show that brine shrimp mortality is inversely related to extract concentrations. *The O. limbata* reported a maximum lethality of 80 % at 150 µg/ml, as seen in the (table no. 6). The highest mortality was reported in 80 % at 150 µg/ml (Fig 5). The bioactive components in both plants make them more cytotoxic than plant extracts.

Table 6. Cytotoxic activity of *Otostegia limbata*

Concentrations µg/ml	Total no Shrimps	Living Shrimps ± SD	Dead Shrimps ± SD	Death %
OLLEE 50 µg/ml	10	4 ± 0.2	6 ± 0.3	60
OLLEE 100 µg/ml	10	3 ± 0.3	7 ± 0.1	70
OLLEE 150 µg/ml	10	2 ± 0.1	8 ± 0.2	80

Key=OLLEE= *Otostegia limbata* Leaf ethanolic extracts, SD =Standard deviations)



**Figure 5.** Graphical representation of cytotoxic activity of *Otostegia limbata*

#### 4. DISCUSSION

The qualitative and quantitative phytochemicals analysis of ethanolic extract was used for detection of phytochemicals. Most of the phytochemicals (alkaloids, terpenes, coumarins, saponins, cardiac glycosides, phlobatannins, flavonoids, quinone, steroids and tannins) were qualitatively and quantitatively detected. Ethanol observed as a best solvent used for the extraction of different phytochemicals. Our results are agreed with the findings of they compare different solvent for phytochemical extraction and found that ethanol is the best solvent for extraction of different phytochemical (Lezoul *et al.*, 2020). Presence of these biologically active compounds shows the medicinal value of *Otostegia limbata* as these phytochemicals have different medicinal properties. *Otostegia limbata*'s phytochemical analysis found that it contains phenols, glycosides, flavonoids, alkaloids, quinones, carbohydrates, amino acids, terpenoids, and coumarins but not tannins, saponins, or sterols, which are thought to be the phytochemicals that give plants their antimicrobial properties (Anthony *et al.*, 2010). Numerous biological processes, including antibacterial, antioxidant, and inflammatory ones, have been connected to flavonoids. They are also known to be able to suppress cell growth and regulate enzymatic activity. They are well recognised to act as a plant's defence mechanism against encroaching diseases (Oikeh *et al.*, 2020). Tannins bind to proline-rich proteins to create complexes that prevent the creation of proteins in cells. It is recognised that the combined effects of tannins, flavonoids, alkaloids, and saponins can stop pathogen growth. Alkaloids are renowned for their anaesthetic, anti-inflammatory, and cardioprotective effects (Javed *et al.*, 2020). Tannins are significant phenolic substances that are well known for their antibacterial properties. The capacity of tannins to precipitate proteins, block the availability of the substrate to the bacterial cells, directly attack the microbial cells, and restrict the uptake of iron by microorganisms is what gives tannins their specific ability to combat dangerous bacterial diseases (Nwankwo *et al.*, 2014). Coumarin's ability to suppress anticholinesterase is what makes them beneficial for treating Alzheimer's disease. Phlobatanins used to cure treating swelling, new wounds, and lymphatic diseases (Kiani *et al.*, 2019). These chemicals are found in *P. stewartii*, according to phytochemical tests, which suggests that this plant is used to treat a variety of illnesses. The *Otostegia limbata* ethanolic extract show the significant activity against all selected oral bacteria strains Table 3, (*Streptococcus mutans*, *Streptococcus mitis*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Bacillus subtilis*, and *Escherichia coli*), this might be due to the antibacterial compound present in our plant are well extracted by ethanol and thus inhibiting the growth of selected bacteria. Our findings are similar to that coded (Philip and

Mahalakshmi, 2019). Using the disc diffusion method, the antibacterial activity of three medicinal plant extracts, including *Azadirachta indica*, *Melia azedarach*, and *Spilanthes acmella*, was examined against *Streptococcus mutans* and *Staphylococcus aureus*.

In order to combat the bacteria that cause denture plaque, plant extracts may be a safe substitute for dangerous medications and chemicals. Similar outcomes were categorized as well (Saquib *et al.*, 2019). They discovered that the two plants' ethanolic extracts had a growth-inhibiting impact on all four strains of periodontal pathobionts. *Tannerella forsythia* was the target of *C. zeylanicum*'s highest antibacterial activity, whereas *Aggregatibacter actinomycetemcomitans* was the target of *C. zeylanicum*'s lowest antibacterial activity among all the groups under study (MIC = 12.5 3.25 mg/mL, MBC = 75 8.23 mg/mL, respectively). *Candida albicans* is significantly inhibited by ethanol extract, although *Aspergillus flavus* is the fungus that is most resistant to our plant. This action demonstrates the antifungal constituents' presence in *O. limbata* and their absorption by the fungi strains. If *A. flavus* activity is lower than usual, it may be because this strain is resistant to the component found in the chosen plant. Our findings concur with the findings (Mohammed *et al.*, 2019). They discovered that *S. marianum* extract works well against *Candida* species at 400-800 g/mL doses. As a result, it was discovered that the plant's fruit sections might be a natural source of antifungal and antibacterial agents. The ethanolic extract of *O. limbata* showed zone of inhibition against *Aspergillus flavus*, *Aspergillus fumigatus*, *Aspergillus Niger*, *Fusarium solani* and *Candida albicans*.

## 5. CONCLUSIONS

By using qualitative phytochemical screening, it was discovered that *Otostegia limbata* possesses active secondary metabolites such as alkaloids, terpenes, coumarins, saponins, cardiac glycosides, phlobatannins, flavonoids, quinone, steroids, and tannins. According to the results of the antimicrobial assay, *Otostegia limbata* plant ethanol extract is a useful tool for testing new antimicrobial medications for the treatment of oral microorganism-related disorders. From the data taken together, it can be inferred that HPLC is a flexible, repeatable chromatographic method for the quantification of medicinal products. Regarding the quantitative and qualitative estimation of active compounds, it has a wide range of applications in various sectors. At 150 g/ml of concentration, the plant likewise exhibits the highest potential for cytotoxic activity. The high performance liquid chromatography potential of *Otostegia limbata* yields impressive results.

## Acknowledgments

The authors are grateful to the Faculty of Health and Biological Science Hazara University Mansehra, Pakistan for providing the facilities for phytochemical analysis, Antimicrobial activity, cytotoxic activity analysis and funding support for this research.

## Conflicts of interest

There is no conflict of interest.

## Data availability statement

All required data is provided in the manuscript; no external dataset is required or utilized.

## REFERENCES

1. Anand U, Jacobo-Herrera N, Altemimi A, Lakhssassi N (2019) A comprehensive review on medicinal plants as antimicrobial therapeutics: potential avenues of biocompatible drug discovery. *Metabolites* **1**, 9, 11, 258.
2. Abbasi AM, Khan MA, Ahmad M, Zafar M, Jahan S, Sultana S (2010) Ethnopharmacological application of medicinal plants to cure skin diseases and in folk cosmetics among the tribal communities of North-West Frontier Province, Pakistan. *Journal of ethnopharmacology* **24**, 128, 2, 322-35.

3. Aga H, Shibuya T, Sugimoto T, Kurimoto M, Nakajima S (1994) Isolation and identification of antimicrobial compounds in Brazilian propolis. *Bioscience, biotechnology, and biochemistry* **1**, 58, 5, 945-6.
4. Anthony F, Diniz LE, Combes MC, Lashermes P (2010) Adaptive radiation in *Coffea* subgenus *Coffea* L.(Rubiaceae) in Africa and Madagascar. *Plant Systematics and Evolution* **285**, 51-64.
5. Arora K, Arora R (2021) Phytochemical Screening, Antioxidant Activity and Antimicrobial Susceptibility of *Trachyspermum ammi* and *Trigonella foenum graecum* Against Dental Caries Causing Microbes.
6. Bankova V, Christoy R, Stoev G, Popov S (1992) Determination of phenolics from propolis by capillary gas chromatography. *Journal of Chromatography* **21**, 607, 1, 150-3.
7. Carron R.A, Maran J.M, Montero L, Fernandozaigo, Dominguez A.A (1987) *Plantas Med. Phytotherapic* **21**, 195-202.
8. Chung KT, Wei CI, Johnson MG (1998) Are tannins a double-edged sword in biology and health? *Trends in Food Science & Technology* **1**, 9, 4,168-75.
9. Drasar PB, Khripach VA (2019) Growing importance of natural products research. *Molecules* **18**, 25, 1:6.
10. Essawi T, Srour M (2000) Screening of some Palestinian medicinal plants for antibacterial activity. *Journal of ethnopharmacology* **15**, 70, 3, 343-9.
11. Enjalric F, Carron MP, Lardet L (1987) Contamination of primary cultures in tropical areas: The case of *Hevea brasiliensis*. Bacterial and Bacteria-like Contaminants of Plant Tissue Cultures **225**, 23, 57-66.
12. Farrukh M, Saleem U, Qasim M, Manan M, Shah MA (2022) *Sarcococca saligna* extract attenuates formaldehyde-induced arthritis in Wistar rats via modulation of pro-inflammatory and inflammatory biomarkers. *Inflammopharmacology* **30**, 2, 579-97.
13. Gamboe RW, Chavez P, Baker WP (2008) Antimicrobial activity of natural plant extracts against five bacterial species. *Journal of the Arizona-Nevada Academy of Science* **40**, 2:165-7.
14. Haq MN, Wazir SM, Ullah F, Khan RA, Shah MS, Khatak A (2016) Phytochemical and biological evaluation of defatted seeds of *Jatropha curcas*. *Sains Malays* **1**, 45(10):1435-42.
15. Hedge I.C, Ali S.I, Nasir Y. J (Eds.) (1990) Labiatae Flora of Pak. (Eds.) No. 192. Karachi & Islamabad.
16. Jamshed H, Siddiqi HS, Gilani AU, Arslan J, Qasim M, Gul B (2019) Studies on antioxidant, hepatoprotective, and vasculoprotective potential of *Viola odorata* and *Wrightia tinctoria*. *Phytotherapy Research* **33**, 9, 2310-8.
17. Javed, B., Nawaz, K. and Munazir, M (2020) Phytochemical analysis and antibacterial activity of tannins extracted from *Salix alba* L. against different gram-positive and gram-negative bacterial strains. *Iranian Journal of Science and Technology*, Transactions A: Science 44, 5,1303-1314.
18. Kale AA, Gaikwada SA, Kamble GS (2011) In vitro anthelmintic activity of stem bark of *Juglans regia* L. *Journal of Chemical and Pharmaceutical Research* **3**, 2, 298-302.
19. Kiani A, Jalili-baleh L, Abdollahi Z, Nadri H, Foroumadi A (2019) Sadat Ebrahimi SE, Khoobi M. Cholinesterase inhibition activity and docking simulation study of coumarin mannich base derivatives. *Journal of Sciences, Islamic Republic of Iran* **26**,30, 1, 5-12.
20. Koo H, Gomes BP, Rosalen PL, Ambrosano GM, Park YK, Cury JA (2000) In vitro antimicrobial activity of propolis and *Arnica montana* against oral pathogens. *Archives of oral biology* **1**, 45, 2, 141-8.
21. Lezoul NE, Belkadi M, Habibi F, Guillén F (2020) Extraction processes with several solvents on total bioactive compounds in different organs of three medicinal plants. *Molecules* **13**, 25, 20, 4672.
22. Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nichols DE, McLaughlin JL (1982) Brine shrimp: a convenient general bioassay for active plant constituents. *Planta medica* **45**, 05:31-4.

23. Mohammed FS, Pehlivan M, Sevindik M (2019) Antioxidant, antibacterial and antifungal activities of different extracts of *Silybum marianum* collected from Duhok (Iraq). *International Journal of Secondary Metabolite* **6**, 4, 317-22.
24. Muhammad H, Qasim M, Ikram A, Versiani MA, Tahiri IA, Yasmeen K, Abbasi MW, Azeem M, Ali ST, Gul B (2020) Antioxidant and antimicrobial activities of *Ixora coccinea* root and quantification of phenolic compounds using HPLC. *South African Journal of Botany* **1**, 135, 71-9.
25. Morrison J, Coromilas JA, Robbins M, Ong LA, Eisenberg S, Stechel RI, Zema MI, Reiser PE, Scherr LA (1980) Digitalis and myocardial infarction in man. *Circulation* **62**, 1, 8-16.
26. Nwankwo, I.U., Onwuakor, C.E. and Aninweze, O.N (2014) Antibacterial activity of Ethanolic extracts of *Citrus sinensis* peels on *Staphylococcus aureus*, *Escherichia coli* and *Pseudomonas aeruginosa* isolated from wound infections. *International Journal of Advances in Pharmacy, Biology and Chemistry* **3**, 4, 941-947.
27. Oikeh EI, Oviasogie FE, Omoregie ES (2020) Quantitative phytochemical analysis and antimicrobial activities of fresh and dry ethanol extracts of *Citrus sinensis* (L.) Osbeck (sweet Orange) peels. *Clinical Phytoscience* **6**, 1, 1-6.
28. Philip JM, Mahalakshmi K (2019) Antimicrobial effect of three Indian medicinal plant extracts on common denture plaque bacteria. *Drug Invention Today* **1**, 11(3).
29. Qasim M, Abideen Z, Adnan MY, Gulzar S, Gul B, Rasheed M, Khan MA (2017) Antioxidant properties, phenolic composition, bioactive compounds and nutritive value of medicinal halophytes commonly used as herbal teas. *South African Journal of Botany* **1**, 110,240-50.
30. Rashid S, Ahmad M, Zafar M, Sultana S, Ayub M, Khan MA, Yaseen G (2015) Ethnobotanical survey of medicinally important shrubs and trees of Himalayan region of Azad Jammu and Kashmir, *Pakistan. Journal of Ethnopharmacology* **26**, 166,340-51.
31. Rosselli S, Fontana G, Bruno M (2019) A review of the phytochemistry, traditional uses, and biological activities of the genus *Ballota* and *Otostegia*. *Planta medica* **85**, 11/12, 869-910.
32. Sadaf HM, Bibi Y, Riaz I, Sultan MA, Bibi F, Bibi M, Hussain M, Sabir S (2016) Pharmacological aptitude and profiling of active constituent from *Otostegia limbata*-Comprehensive review. *Asian Pacific Journal of Tropical Disease* **1**, 6, 11, 918-24.
33. Saquib SA, AlQahtani NA, Ahmad I, Kader MA, Al Shahrani SS, Asiri EA (2019) Evaluation and comparison of antibacterial efficacy of herbal extracts in combination with antibiotics on periodontal pathobionts: an in vitro microbiological study. *Antibiotics* **1**, 8, 3, 89.
34. Scheen AC, Albert VA (2007) Nomenclatural and taxonomic changes within the *Leucas* clade (*Lamioideae*; *Lamiaceae*). *Systematics and Geography of Plants* **1**, 229-38.
35. World Health Organization. The world health report (2002) reducing risks, promoting healthy life. World Health Organization.