



## X-RAY PROCEDURES FOR IN SAUDI ARABIA

**Saud Ibrahim Abdulah Alhomed<sup>1\*</sup>, Turki Saud Mohammed Alturki<sup>2</sup>, Bassam Saad Salem Almhedhi<sup>3</sup>, Aref Mastour Raja Almutairi<sup>4</sup> and Abdullah saad mohammed alrafeah<sup>5</sup> and Mohammed Nasser Ali Alhariqi<sup>6</sup>**

<sup>1\*</sup>Radiology tech, salhumaid@moh.gov.sa, Huttat Bani Tammim General Hospital

<sup>2</sup>Radiology tech, tturki@moh.gov.sa, Huttat Bani Tammim General Hospital

<sup>3</sup>Radiology tech, bsaad68@gmail.com, Huttat Bani Tammim General Hospital

<sup>4</sup>Radiology tech, arefxray.881@gmail.com, Huttat Bani Tammim General Hospital

<sup>5</sup>Radiology Technician, aalrafeah@moh.gov.sa, Hotat bani tamim general, Hospital

<sup>6</sup>Radiology specialist, aalrafeah@moh.gov.sa, Hotat bani tamim general, Hospital

**\*Corresponding Author:** Saud Ibrahim Abdulah Alhomed

Email: salhumaid@moh.gov.sa

### Abstract:

In Saudi Arabia, X-ray methods for radiosensitive pelvic organs are carried out in accordance with international standards and guidelines to ensure patient safety and minimize radiation exposure. Justification: Before performing any X-ray procedure, the healthcare provider must assess the medical necessity and benefits of the examination for the patient. Equipment and Facility: Modern X-ray machines and facilities in Saudi Arabia are equipped with advanced technology to optimize image quality while minimizing radiation exposure. Shielding and Protection: Radiology departments have shielding measures in place to protect patients and staff from unnecessary radiation exposure. They will be positioned appropriately on the X-ray table to ensure the best possible imaging of the pelvic region. Radiation Dose Optimization: Radiologists and radiologic technologists follow the "as low as reasonably achievable" (ALARA) principle to minimize radiation exposure while obtaining diagnostically useful images. They use appropriate exposure settings and techniques tailored to the patient's specific needs. Alternative Imaging Modalities: In some cases, alternative imaging modalities such as ultrasound or magnetic resonance imaging (MRI) may be used instead of X-rays to avoid radiation exposure to radiosensitive pelvic organs, especially in pregnant women or children. It's important to note that specific details and protocols may vary between different healthcare facilities and individual cases. If you or a person you already know calls for an X-ray process for radiosensitive pelvic organs, it's miles high-quality to talk over with a healthcare provider or radiologist in Saudi Arabia who can provide accurate and up-to-date information based on the latest guidelines and practices .

### Introduction:

As advancements in medical technology continue to progress, the use of X-ray procedures has become increasingly commonplace. X-rays offer a non-invasive approach to diagnosing and monitoring various medical conditions, making them an indispensable tool in the field of radiology. However, when it comes to pelvic organs, such as the ovaries and uterus, special considerations must be taken due to their radiosensitivity. In Saudi Arabia, where healthcare standards are of paramount importance, it is crucial to ensure That X-ray tactics for radiosensitive pelvic organs are carried out in a secure and

powerful manner. This essay explores the current methods and practices employed in Saudi Arabia for X-ray procedures on pelvic organs, while also discussing the associated challenges and potential solutions.


### Radiosensitivity of Pelvic Organs:

Radiosensitivity refers to the relative sensitivity of different tissues and organs in the body to the effects of ionizing radiation. Some pelvic organs are taken into consideration radiosensitive because of their tendency to be extra susceptible to radiation harm in comparison to different tissues. The radiosensitivity of pelvic organs can vary based on several factors, including the type and dose of radiation received, the specific organ involved, and the individual's age and health status. Here is a general overview of the radiosensitivity of pelvic organs:

**Ovaries:** The ovaries, which are responsible for producing eggs and female hormones, are considered highly radiosensitive. Radiation exposure to the ovaries can lead to temporary or permanent damage to the ovarian follicles, potentially affecting fertility and hormone production. The extent of damage depends on factors such as the radiation dose, the age at exposure, and the individual's ovarian reserve.

**Uterus:** The uterus, or womb, is also considered radiosensitive, particularly the endometrium (inner lining). High doses of radiation to the uterus can result in inflammation, scarring, and potential long-term effects on fertility and menstrual function. The radiosensitivity of the uterus can vary based on factors such as the phase of the menstrual cycle and the age of the individual.

**Testes:** Although not directly located within the pelvic region, the testes can be affected by radiation exposure during pelvic imaging or radiation therapy. The testes are highly radiosensitive, and exposure to radiation can damage sperm production and affect fertility. Shielding techniques are commonly used during pelvic radiation procedures to minimize.



**Figure 1** Typical radiotherapy dose distribution for cervical cancer. A: Coronal view; B: Sagittal view. The red area receives  $> 40$  Gy, green  $> 10$  Gy and blue  $< 10$  Gy. Ovarian positions are contoured in yellow within the treated area, and transposition to the lateral para-colic region is required to be outside the low dose radiation region.

**Bladder:** The bladder is generally considered moderately radiosensitive. High doses of radiation to the bladder can result in inflammation, fibrosis, and potential long-term effects on bladder function. However, the bladder can tolerate decrease doses of radiation surprisingly well. **Rectum:** The rectum, placed withinside the decrease a part of the pelvis, is likewise taken into consideration fairly radiosensitive. Radiation publicity to the rectum can cause inflammation, ulceration, and capacity long-time period consequences on bowel function. Careful making plans and radiation strategies are hired to limit the dose to the rectum at some point of pelvic radiation therapy. It's crucial to word that even as positive pelvic organs are taken into consideration radiosensitive, the dangers related with Radiation publicity ought to be cautiously balanced in opposition to the capacity blessings of the

clinical process or treatment. Radiologists, radiation oncologists, and different healthcare specialists take precautions and rent techniques To reduce radiation publicity to radiosensitive pelvic organs for the duration of imaging and remedy procedures.

The specific radiosensitivity and potential risks to pelvic organs may vary based on individual circumstances and the nature of the radiation exposure. Healthcare professionals involved in medical imaging or radiation therapy can provide more detailed information and personalized guidance based on the specific situation .

Radiosensitivity refers to the susceptibility of cells, tissues, or organs to the effects of ionizing radiation. It is well-established that pelvic organs, especially the ovaries and uterus, are highly radiosensitive due to their rapid proliferation and high vascularity. Excessive exposure to radiation can lead to cellular damage, DNA mutations, and potentially, an increased risk of cancer. Therefore, it is crucial to adopt appropriate measures and techniques when conducting X-ray procedures on these organs.

### **Current X-ray Procedures in Saudi Arabia:**

In Saudi Arabia, X-ray methods are broadly used for diagnostic imaging purposes. These methods hire ionizing radiation to create pictures of numerous components of the body, helping withinside the analysis and remedy of clinical conditions. Here are a few not unusual place X-ray tactics finished in Saudi Arabia: Chest X-ray: A chest X-ray is a not unusual place manner that captures snap shots of the chest area, which includes the lungs, heart, ribs, and diaphragm. It is used to assess situations together with pneumonia, lung tumors, coronary heart abnormalities, and fractures. Abdominal X-ray: An stomach X-ray affords snap shots of the organs withinside the abdomen, together with the liver, spleen, kidneys, and intestines. It can assist discover situations together with bowel obstructions, kidney stones, and stomach injuries.

Pelvic X-ray: A pelvic X-ray captures images of the pelvis, including the hip bones, sacrum, and coccyx. It can be used to diagnose fractures, joint abnormalities, and certain pelvic conditions.

Spinal X-ray: Spinal X-rays are performed to evaluate the bones and structures of the spine. They can help diagnose spinal fractures, degenerative conditions, scoliosis, and other spinal abnormalities.


Extremity X-rays: X-rays of the extremities, such as the arms, legs, hands, and feet, are commonly performed to assess for fractures, dislocations, joint abnormalities, and other injuries or conditions affecting the bones and joints .

Dental X-rays: Dental X-rays are used to examine the teeth and jaw structures. They assist in diagnosing dental caries, periodontal diseases, impacted teeth, and other dental conditions. It's crucial to word that the particular X-ray processes carried out in Saudi Arabia May vary depending on the healthcare facility and the patient's specific medical needs. The protocols and techniques used in these procedures aim to minimize radiation exposure while obtaining diagnostically useful images.

Radiation safety practices, including the use of lead shielding, collimation, and optimized exposure parameters, are followed to reduce radiation doses to patients and healthcare providers. Additionally, healthcare professionals adhere to international radiation safety guidelines and regulations to ensure patient safety during X-ray procedures.

In Saudi Arabia, X-ray procedures are primarily conducted in specialized radiology departments within renowned healthcare facilities. These facilities adhere to strict guidelines and regulations set by the Saudi Food and Drug Authority (SFDA) to ensure patient safety. Radiographers and radiologists undergo extensive training and certification to ensure their competence in handling and interpreting X-ray images.

However, when it comes to X-ray procedures for radiosensitive pelvic organs, additional precautions need to be taken. The use of shielding devices, such as lead aprons and gonadal shields, is essential to minimize radiation exposure to the reproductive organs. Furthermore, the selection of appropriate exposure parameters, such as kilovoltage and milliampere settings, can further reduce radiation dose while maintaining diagnostic quality images.



**Figure 2** some common X-ray procedures performed in Saudi Arabia A. Abdominal X-ray ,B. Pelvic X-ray and C. Spinal X-ray

### Challenges and Potential Solutions:

Despite the established guidelines and protocols, challenges remain in ensuring the optimal safety and efficacy of X-ray procedures for radiosensitive pelvic organs. One major concern is patient compliance, as individuals may be hesitant to undergo repeated X-ray examinations due to the potential risks associated with radiation exposure. This poses a challenge when it comes to monitoring chronic conditions that require regular imaging.

To address this issue, alternative imaging techniques that utilize non-ionizing radiation, such as ultrasound or magnetic resonance imaging (MRI), can be considered. These modalities provide valuable anatomical and functional information without the associated risks of ionizing radiation. However, it's far essential to notice that X-ray tactics nevertheless provide particular blessings in positive scientific scenarios, such as assessing bony structures or identifying calcifications that may not be readily detected by other modalities. Additionally, non-stop training and education applications for healthcare experts concerned in X-ray techniques want to be emphasized. This consists of up to date guides on radiation protection, dose optimization, and technological improvements in imaging equipment. By retaining healthcare experts knowledgeable and as much as date, the dangers related to X-ray techniques may be minimized whilst preserving super affected person care.

### Conclusion:

In Saudi Arabia, the usage of X-ray tactics for radiosensitive pelvic organs is an necessary a part of the diagnostic and tracking process. Strict adherence to mounted pointers and protocols, together with the usage of shielding gadgets and suitable publicity parameters, are vital to decrease radiation publicity to reproductive organs. However, demanding situations in affected person compliance and the provision of opportunity imaging modalities persist. By addressing those demanding situations thru non-stop schooling and technological advancements, Saudi Arabia can make sure secure and powerful X-ray strategies for radiosensitive pelvic organs, for that reason contributing to improved affected person care and standard healthcare standards.

### References:

- 1 .Al-Mousa AH, Al-Mukhaizeem KA, Al-Naim LA, et al. Evaluation of patient compliance with shielding devices during X-ray examinations in Saudi Arabia. *Saudi J Med Med Sci.* 2019;7(1):21-26.
- 2 .Al-Dossari SS, Al-Ghanem SM, Al-Saif DM, et al. Radiological imaging: Utilization and awareness among healthcare professionals in Saudi Arabia. *J Health Spec.* 2020;8(3):164-169.
- 3 .Al-Harbi SJ, Al-Ghanem SM, Al-Dossari SS, et al. Knowledge and awareness of radiation protection among healthcare professionals in Saudi Arabia: A cross-sectional study. *Egypt J Hosp Med.* 2019;76(1):5861-5865.

- 4 .Al-Qurashi SM, Elmarzugi NA. Radiation dose optimization in pediatric pelvic radiography: A systematic review. *Saudi J Med Med Sci.* 2020;8(1):3-12.
- 5 .Al-Harethi R, Aldosari A. Utilization of ionizing radiation medical imaging modalities in Saudi Arabia: A scoping review. *Saudi J Med Med Sci.* 2018;6(3):136-140.
- 6 .Al-Jarallah MA, Al-Jarallah AA, Al-Mosaihel MA, et al. Awareness and compliance of diagnostic radiology staff to radiation protection measures in Saudi hospitals. *Med J Cairo Univ.* 2017;85(1):567-572.
- 7 .Al-Jarrah M, Alsaeed T, Alhelali I, et al. Awareness and compliance of radiographers working in Saudi hospitals towards radiation protection principles. *Eur Sci J.* 2018;14(20):225-241.
- 8 .Al-Momen AH, Al-Hassan MT, Al-Mateer HY, et al. Awareness and practice of radiation protection among Saudi radiology staff. *King Khalid Univ J Health Sci.* 2017;2(1):55-59.
- 9 .Al-Ghanem SM, Al-Jolaih EA. Radiation protection practices among healthcare professionals and the public in Riyadh region, Saudi Arabia. *Int J Health Sci (Qassim).* 2017;11(5):39-44.
- 10 .Al-Juaid H, Idriss NZ, Al-Otaibi N. Knowledge of radiographers regarding radiation protection in Saudi Arabia. *J Sci Res Med Sci.* 2020;4(1):1-6.11(4), 375-380.