RESEARCH ARTICLE DOI: 10.53555/jptcp.v31i4.5466

# "MORINGA OLEIFERA IN THE TREATMENT OF OSTEOPOROSIS: A COMPREHENSIVE REVIEW OF MECHANISMS, EFFICACY, AND FUTURE DIRECTIONS"

Naruboyina Vasavi<sup>1\*</sup>, L.S. Jyothika<sup>2</sup>, Y. Sai Charan Teja<sup>3</sup>, K. Somasekhar Reddy<sup>4</sup>, D. Yamini Kalyani<sup>5</sup>

1\*,2,3,4,5 Department of Pharmacology, Raghavendra Institute of Pharmaceutical Education and Research (RIPER) – Autonomous, KR Palli cross, Chiyyedu (post), Anantapur, Andhra Pradesh,515721, India.

# \*Corresponding Author: Naruboyina Vasavi

\*Department of Pharmacology, Raghavendra Institute of Pharmaceutical Education and Research (RIPER) – Autonomous, KR Palli cross, Chiyyedu (post), Anantapur, Andhra Pradesh,515721, India.

#### **Abstract:**

Osteoporosis is a pressing health concern characterized by diminished bone density and increased susceptibility to fractures, particularly among the elderly. The condition poses significant challenges due to its silent progression and substantial impact on quality of life and healthcare systems. Addressing osteoporosis requires effective prevention and treatment strategies. Moringa oleifera, commonly known as the "drumstick tree," has garnered attention for its potential therapeutic effects in managing osteoporosis. This review outlines the accumulating evidence suggesting that Moringa oleifera may offer promising avenues for osteoporosis management. Notably, Moringa oleifera contains a plethora of bioactive components, including vitamins, minerals, flavonoids, and glucosinolates, which contribute to its medicinal properties. Exploration of the mechanisms of action underlying Moringa oleifera's effects on bone health reveals its potential to promote bone formation, inhibit bone resorption, and exert anti-inflammatory and antioxidant effects. Preclinical studies have demonstrated encouraging results, showing improvements in bone mineral density, bone architecture, and biomarkers of bone turnover. Furthermore, clinical evidence supports the efficacy and safety of Moringa oleifera supplementation in osteoporosis patients. However, further research is warranted to elucidate optimal dosing regimens, evaluate long-term outcomes, and address potential confounding factors. In conclusion, this review highlights the potential of Moringa oleifera as a natural adjunctive therapy for osteoporosis. By synthesizing current knowledge on its bioactive components, mechanisms of action, preclinical and clinical evidence, and future research directions, this review aims to contribute to the growing understanding of Moringa oleifera's role in osteoporosis management.

#### 1. Introduction

Osteoporosis stands as a prevalent and pressing global health issue, characterized by the progressive weakening of bone density and microarchitecture, leading to an increased risk of fractures. This condition affects millions worldwide, particularly among the elderly and postmenopausal women, posing significant challenges to public health systems and individual well-being. (1) Despite

advancements in medical science, current treatments for osteoporosis, such as bisphosphonates and hormone replacement therapy, have limitations including potential side effects and suboptimal efficacy in some patients. <sup>(2)</sup> Consequently, there is a growing recognition of the need for alternative or adjunctive therapies to complement existing treatments and improve outcomes for individuals with osteoporosis<sup>(3)</sup>

In recent years, attention has turned to natural remedies as potential options for managing osteoporosis. One such remedy that has gained increasing interest is Moringa oleifera, a plant native to the Indian subcontinent and widely cultivated in tropical and subtropical regions worldwide. Moringa oleifera, also known as the "drumstick tree" or "miracle tree," has a long history of traditional medicinal use for various health conditions due to its rich nutritional and pharmacological properties. (4) Moringa oleifera is renowned for its exceptional nutritional profile, containing high levels of vitamins, minerals, proteins, and antioxidants. (5) Additionally, it is abundant in bioactive compounds such as flavonoids, phenolic acids, and glucosinolates, which contribute to its diverse therapeutic effects. (6) Emerging research suggests that these bioactive components may hold promise in mitigating the progression of osteoporosis and promoting bone health.

Studies investigating the potential effects of Moringa oleifera on bone metabolism have shown promising results in preclinical models. Animal studies have demonstrated improvements in bone mineral density, bone microarchitecture, and biomarkers of bone turnover following supplementation with Moringa oleifera extracts. <sup>(7)</sup> These findings have spurred interest in further exploring the mechanisms of action underlying its effects on bone health. In light of the limitations of current osteoporosis treatments and the promising preliminary evidence, there is a growing interest in investigating the therapeutic potential of Moringa oleifera as a natural remedy for osteoporosis. This review aims to provide a comprehensive overview of the bioactive components, mechanisms of action, preclinical and clinical evidence, and future research directions pertaining to the use of Moringa oleifera in osteoporosis management. <sup>(8)(9)</sup>

#### 2. Osteoporosis in Menopausal Women:

Hormonal changes during menopause have a profound impact on bone health. The key hormonal changes that occur during menopause include a significant decline in estrogen and changes in other hormones like progesterone and parathyroid hormone. These changes lead to several consequences that affect bone health.

#### 1. Estrogen Decline:

Estrogen plays a crucial role in maintaining bone density by regulating the balance between bone formation and bone resorption. During menopause, there is a sharp decrease in estrogen levels. This hormonal shift leads to accelerated bone resorption (the process of breaking down bone tissue) and a reduction in bone formation. As a result, bone density decreases more rapidly. (10)

#### 2. Loss of Bone Mass:

Menopausal women often experience a loss of bone mass, particularly in the trabecular bone, which is the spongy interior of bones. This loss of bone mass compromises bone strength and increases the likelihood of fractures. (11)

#### 3. Increased Bone Porosity:

As bone density diminishes, bones become more porous and less dense. This increased porosity weakens bone structure and reduces resistance to fractures. (12)

#### 4. Higher Fracture Risk:

The combination of reduced bone density, increased bone porosity, and the loss of bone mass makes menopausal women more susceptible to fractures. Common sites for osteoporosis-related fractures during this stage include the hip, spine, and wrist. (13)

#### 5. Impact on Bone Remodelling:

Hormonal changes during menopause disrupt the natural bone remodelling process. This process involves the removal of old bone tissue and the formation of new bone. With the estrogen decline, this balance is skewed, resulting in more bone being broken down than rebuilt. (14)

## 6. Quality of Life and Independence:

Fractures, particularly those involving the hip or spine, can have profound consequences for a woman's overall quality of life. They often lead to physical pain, decreased mobility, and loss of independence. Women may experience limitations in their ability to perform daily activities, leading to a decrease in overall well-being.

#### 7. Financial and Emotional Burdens:

Managing fractures and their aftermath can lead to significant financial and emotional burdens. This includes healthcare costs, rehabilitation, and caregiving. These burdens can affect a woman's emotional well-being and that of her family. (15)

## 2.1 Mechanism of estrogen decline in menopause.

The mechanism of estrogen decline in menopause is primarily associated with the aging of a woman's reproductive system. Estrogen, produced primarily in the ovaries, plays a pivotal role in regulating the female reproductive system and maintaining various bodily functions. During menopause, several factors contribute to the progressive decrease in estrogen levels:

#### 1. Ovarian Aging:

The primary source of estrogen in premenopausal women is the ovaries. Ovarian aging is a natural process that occurs throughout a woman's life. As a woman ages, the number of follicles (tiny sacs that contain immature eggs) in her ovaries diminishes. With fewer follicles, there is a reduced capacity to produce and release estrogen.

#### 2. Follicular Atresia:

As women age, many of their ovarian follicles undergo atresia, which is the degeneration and reabsorption of these follicles. This process further reduces the number of functional follicles and, consequently, the estrogen-producing capacity of the ovaries.

## 3. Menstrual Irregularities:

As the number of functional follicles decreases, menstrual cycles may become irregular. Ovulation may not occur with each cycle, leading to variations in estrogen production.

## 4. Aging Ovarian Follicles:

The remaining ovarian follicles may produce less estrogen as they age, and the quality of the eggs within them declines. This contributes to a decrease in estrogen levels and an overall reduction in reproductive capacity.

#### 5. Gradual Hormonal Shift:

As the ovarian function declines, there is a shift in hormonal balance. The hypothalamus and pituitary gland in the brain, which regulate reproductive hormones, attempt to compensate for the declining ovarian function by releasing higher levels of follicle-stimulating hormone (FSH) and luteinizing hormone (LH). These hormones stimulate the ovaries to produce more estrogen, but the ovaries' capacity to respond diminishes.

## 6. Menopause Onset:

Eventually, the ovaries reach a point where they can no longer respond to FSH and LH effectively. This results in a sharp decline in estrogen production, marking the onset of menopause.

The process of estrogen decline is gradual, and it typically occurs over several years leading up to menopause, known as perimenopause. During perimenopause, women may experience irregular menstrual cycles, hot flashes, night sweats, and other symptoms related to hormonal fluctuations. Once menopause is reached, estrogen levels remain consistently low. (16)

It's important to note that while the ovaries are the primary source of estrogen, other tissues in the body, such as fat cells and the adrenal glands, continue to produce small amounts of estrogen even

after menopause. However, these sources are not sufficient to maintain premenopausal levels of estrogen, leading to the characteristic hormonal changes associated with menopause.

#### 2.2 Mechanism of loss of bone mass in osteoporosis

The loss of bone mass in osteoporosis is a complex process that involves an imbalance between bone formation and bone resorption. Several cellular and molecular mechanisms contribute to this condition.

#### 1. Normal Bone Remodelling:

Bone tissue undergoes a continuous remodelling process, which involves the removal of old bone (resorption) and the formation of new bone (formation). This process helps maintain bone strength and repair microdamage.

## 2. Osteoclast Activity:

Osteoclasts are specialized cells responsible for bone resorption. They secrete enzymes and acids that break down the mineralized bone matrix and release calcium into the bloodstream. This is a normal part of bone remodelling.

#### 3. Osteoblast Activity:

Osteoblasts are cells responsible for bone formation. They produce collagen and other proteins that make up the bone matrix and initiate the process of mineralization, which strengthens bone.

#### 4. Hormonal Regulation:

Hormones play a crucial role in maintaining the balance between bone formation and resorption. Estrogen has a significant impact on bone health. It suppresses the activity of osteoclasts, reducing bone resorption and promoting bone formation.

## 5. The Role of Parathyroid Hormone (PTH):

Parathyroid hormone is another hormone that regulates calcium and phosphate levels in the body. In response to low blood calcium levels, the parathyroid glands release PTH, which stimulates osteoclast activity, leading to increased bone resorption to release calcium. (17)

#### 6. Bone Loss in Osteoporosis:

Osteoporosis occurs when there is an imbalance in bone remodelling, with increased bone resorption and decreased bone formation. Several factors contribute to this imbalance:

- Hormonal Changes: In postmenopausal women, the significant decline in estrogen levels (as discussed earlier) results in an increase in osteoclast activity and bone resorption, leading to bone loss.
- Aging: Aging itself is associated with changes in bone density. As individuals get older, bone turnover slows down, and the balance between resorption and formation becomes skewed, with a greater emphasis on resorption.
- Nutritional Deficiencies: Inadequate intake of essential nutrients like calcium and vitamin D can compromise bone health and reduce the ability to form new bone tissue.
- Lifestyle Factors: Sedentary lifestyles, smoking, excessive alcohol consumption, and certain medications can further accelerate bone loss.

#### 7. Microarchitectural Changes:

In osteoporosis, the loss of bone mass is not uniform. Instead, there are microarchitectural changes in bone structure, making the bone more porous and less dense. This microarchitectural deterioration weakens the bone and increases its fragility.

#### 8. Fracture Risk:

Osteoporosis results in bones that are more susceptible to fractures, particularly in weight-bearing areas like the hip, spine, and wrist. These fractures can have a profound impact on an individual's overall quality of life, independence, and well-being. (18)

#### 2.3 Mechanism of Increased Bone Porosity

Increased bone porosity, also known as osteoporosis, is a condition characterized by the deterioration of bone microarchitecture, leading to decreased bone density and increased

vulnerability to fractures. Several mechanisms contribute to the development of increased bone porosity:

#### 1. Enhanced Bone Resorption:

Osteoclasts, specialized cells responsible for breaking down bone tissue, play a central role in bone remodelling. In osteoporosis, there is an imbalance between bone resorption and formation, with increased activity of osteoclasts. This excessive bone resorption results in the removal of bone matrix, leading to the creation of larger and more numerous pores within the bone structure.

#### 2. Reduced Bone Formation:

Osteoblasts, cells responsible for bone formation, are also affected in osteoporosis. While bone formation occurs concurrently with resorption during normal bone remodelling, the activity of osteoblasts may be insufficient to compensate for the excessive bone resorption in osteoporosis. This leads to a net loss of bone mass and increased porosity.

#### 3. Hormonal Imbalance:

Hormonal changes, particularly the decline in estrogen levels in postmenopausal women, contribute to increased bone porosity. Estrogen plays a crucial role in maintaining bone density by inhibiting osteoclast activity and promoting osteoblast function. The decrease in estrogen levels during menopause disrupts this balance, leading to accelerated bone loss and increased porosity.

## 4. Age-Related Changes:

Aging is a significant risk factor for osteoporosis. As individuals age, bone remodelling becomes less efficient, resulting in decreased bone formation and increased bone resorption. This age-related decline in bone turnover contributes to the gradual loss of bone mass and increased porosity.

#### 5. Nutritional Deficiencies:

Inadequate intake of essential nutrients such as calcium, vitamin D, and protein can impair bone health and contribute to increased bone porosity. Calcium and vitamin D are essential for bone mineralization and remodelling processes, while protein provides the building blocks necessary for bone formation.

#### 6. Lifestyle Factors:

Certain lifestyle factors, including sedentary behaviour, smoking, excessive alcohol consumption, and poor dietary habits, can exacerbate bone loss and increase porosity. These factors may directly impact bone turnover or indirectly affect bone health through their effects on hormonal regulation, nutrient absorption, and overall bone metabolism. (19)

#### 7. Microarchitectural Changes:

The structural integrity of bone is compromised in osteoporosis, leading to alterations in bone microarchitecture. These microarchitectural changes include thinner trabeculae (the lattice-like structures within bone), increased trabecular separation, and larger intertrabecular spaces. These alterations result in decreased bone density and increased porosity, making the bone more susceptible to fractures.

#### 2.4 Higher Fracture Risk

Higher fracture risk is a significant consequence of osteoporosis, a condition characterized by reduced bone density and increased bone fragility. Several factors contribute to the elevated risk of fractures in individuals with osteoporosis.

#### 1. Decreased Bone Strength:

Osteoporosis leads to the deterioration of bone microarchitecture, resulting in bones that are less dense and more porous. This structural weakness compromises bone strength and resistance to mechanical stress, making fractures more likely.

## 2. Loss of Bone Mass:

Osteoporosis is associated with a decrease in bone mass, particularly in trabecular bone (the spongy inner layer of bone) but also in cortical bone (the dense outer layer). The loss of bone mass weakens the structural integrity of the bone, increasing susceptibility to fractures. (20)

#### 3. Site-Specific Fracture Patterns:

Osteoporosis-related fractures often occur at specific sites, including the hip, spine (vertebrae), and wrist. Hip fractures are particularly concerning due to their association with significant morbidity, mortality, and loss of independence. Vertebral fractures can lead to pain, spinal deformities, and height loss, while wrist fractures can impair hand function and mobility.

## 4. Age-Related Changes:

Aging is a major risk factor for osteoporosis and fractures. As individuals age, bone turnover decreases, leading to gradual bone loss and decreased bone strength. Age-related changes in bone structure, hormonal regulation, and muscle mass further contribute to the increased fracture risk in older adults. (21)

#### 5. Hormonal Factors:

Hormonal changes play a critical role in bone metabolism, and alterations in hormone levels can affect bone health. Postmenopausal women are particularly susceptible to osteoporosis and fractures due to the decline in estrogen levels following menopause. Estrogen plays a key role in maintaining bone density by inhibiting bone resorption and promoting bone formation. Similarly, low levels of testosterone in men can also contribute to bone loss and increased fracture risk.

#### 6. Nutritional Deficiencies:

Inadequate intake of essential nutrients such as calcium, vitamin D, and protein can weaken bones and increase fracture risk. Calcium and vitamin D are essential for bone mineralization and remodelling processes, while protein provides the building blocks necessary for bone formation. (23)

## 7. Lifestyle Factors:

Certain lifestyle factors can exacerbate bone loss and increase fracture risk. Sedentary behaviour, smoking, excessive alcohol consumption, and poor dietary habits can negatively impact bone health. Additionally, falls resulting from impaired balance, muscle weakness, or environmental hazards further increase the likelihood of fractures in individuals with osteoporosis.

#### 8. Previous Fractures:

Individuals who have already experienced a fracture due to osteoporosis are at increased risk of subsequent fractures. This increased risk may be attributed to underlying bone fragility and compromised bone healing mechanisms.

#### 2.5 Impact on Bone Remodelling

Bone remodelling is a continuous process that involves the removal of old bone tissue (resorption) by osteoclasts and the formation of new bone tissue (formation) by osteoblasts. This dynamic process helps maintain bone strength, repair microdamage, and regulate mineral homeostasis. However, in conditions like osteoporosis, there is an imbalance between bone resorption and formation, leading to a disruption in bone remodelling.

#### 1. Increased Bone Resorption:

Osteoclasts, specialized cells responsible for bone resorption, are overactive in osteoporosis. They secrete enzymes and acids that break down the mineralized bone matrix, leading to the removal of old bone tissue. This excessive bone resorption contributes to the loss of bone mass and structural deterioration characteristic of osteoporosis.

#### 2. Reduced Bone Formation:

While bone formation occurs concurrently with resorption during normal bone remodelling, the activity of osteoblasts, cells responsible for bone formation, may be insufficient to compensate for the increased bone resorption in osteoporosis. This results in a net loss of bone mass over time. Factors contributing to reduced bone formation in osteoporosis include hormonal imbalances, agerelated changes, and nutritional deficiencies.

## 3. Imbalance in Remodelling Cycle:

In osteoporosis, the balance between bone resorption and formation is disrupted, leading to an imbalance in the remodelling cycle. The duration of bone resorption may be prolonged, while the duration of bone formation may be shortened. This altered remodelling cycle results in a negative bone turnover balance, with more bone being removed than replaced.

## 4. Changes in Bone Microarchitecture:

Osteoporosis is characterized by alterations in bone microarchitecture, including thinner trabeculae (the lattice-like structures within bone), increased trabecular separation, and larger intertrabecular spaces. These microarchitectural changes weaken bone structure and reduce bone strength, making bones more susceptible to fractures.

#### 5. Site-Specific Effects:

Osteoporosis affects bone remodelling differently in various skeletal sites. Trabecular bone, which is found in the spine and ends of long bones, is more susceptible to osteoporotic changes than cortical bone, which forms the dense outer layer of long bones. As a result, osteoporosis-related fractures often occur in trabecular-rich areas like the spine, hip, and wrist.

## 6. Impaired Bone Healing:

Osteoporosis can impair the bone healing process following a fracture. The compromised bone microenvironment, reduced bone formation capacity, and altered cellular activity in osteoporotic bone may hinder the normal bone healing response, leading to delayed healing, non-union, or malunion of fractures. (23)

#### 2.5 Quality of Life and Independence

The quality of life and independence of individuals with osteoporosis can be significantly impacted due to the physical, psychological, and social consequences of the condition. (24)

#### 1. Physical Impairment:

Pain: Osteoporosis-related fractures, especially those in the spine and hip, can cause significant pain, reducing mobility and impairing daily activities.

Decreased Mobility: Fractures, pain, and skeletal deformities resulting from osteoporosis can limit mobility, making it difficult to walk, climb stairs, or perform routine tasks independently.

Loss of Height: Vertebral fractures can lead to compression of the spine, resulting in a reduction in height. This can affect posture and increase the risk of falls.

Impaired Physical Function: Fractures and skeletal deformities can impair physical function, affecting activities such as dressing, bathing, and household chores.

#### 2. Psychological Impact:

Anxiety and Depression: Chronic pain, functional limitations, and concerns about future fractures can contribute to anxiety and depression in individuals with osteoporosis.

Body Image Issues: Changes in posture and height due to vertebral fractures can impact body image and self-esteem, leading to feelings of self-consciousness and social withdrawal.

Fear of Falling: Individuals with osteoporosis may develop a fear of falling, which can further restrict activity levels and reduce quality of life.

#### 3. Social Consequences:

Loss of Independence: Functional limitations and mobility impairments can lead to a loss of independence, as individuals may rely on others for assistance with daily tasks and activities.

Social Isolation: Pain, fear of falling, and mobility restrictions can result in social isolation and withdrawal from social activities and interactions.

Financial Burden: Osteoporosis-related fractures can impose a financial burden on individuals and their families due to healthcare costs, including medical treatments, rehabilitation, and assistive devices.

#### 4. Impact on Care givers:

Increased Caregiver Burden: Family members or caregivers may experience increased stress and burden due to the caregiving responsibilities associated with supporting individuals with osteoporosis, especially those with mobility limitations or cognitive impairment.

Emotional Distress: Caregivers may experience emotional distress, including feelings of guilt, frustration, and burnout, as they navigate the challenges of supporting a loved one with osteoporosis.

#### 5. Reduced Quality of Life:

Overall, osteoporosis can significantly reduce the quality of life for individuals affected by the condition, impacting physical function, psychological well-being, social engagement, and overall satisfaction with life.

#### 2.6 Financial and Emotional Burdens

The financial and emotional burdens associated with osteoporosis can significantly impact individuals, families, and healthcare systems.

#### 1. Healthcare Costs:

Medical Treatments: Osteoporosis-related healthcare costs include expenses associated with diagnostic tests (e.g., bone density scans), medications (e.g., bisphosphonates, calcium, vitamin D supplements), and surgical interventions (e.g., fracture repair, joint replacement).

Rehabilitation: Individuals with osteoporosis may require rehabilitation services, such as physical therapy and occupational therapy, to recover from fractures and regain functional independence.

Assistive Devices: Mobility aids (e.g., walkers, canes), orthotic devices (e.g., back braces), and home modifications (e.g., handrails, grab bars) may be necessary to support individuals with osteoporosis and prevent falls.

#### 2. Lost Productivity:

Work Absenteeism: Osteoporosis-related pain, fractures, and medical appointments may result in missed workdays or reduced work productivity for individuals affected by the condition.

Caregiver Responsibilities: Family members or caregivers may need to take time off work to provide care and support for individuals with osteoporosis, resulting in lost wages and productivity.

#### 3. Out-of-Pocket Expenses:

Co-payments and Deductibles: Individuals with osteoporosis may incur out-of-pocket expenses for insurance co-payments, deductibles, and uncovered healthcare services.

Over-the-Counter Medications: Over-the-counter pain relievers and supplements may be necessary to manage osteoporosis-related symptoms and complications, adding to out-of-pocket costs. (25)

#### 4. Long-Term Care:

Nursing Home Care: Severe osteoporosis-related fractures or complications may require long-term nursing home care, which can be financially burdensome for individuals and their families.

In-Home Care: Some individuals with osteoporosis may require in-home care services, such as assistance with activities of daily living (e.g., bathing, dressing), meal preparation, and medication management.

#### 5. Emotional impact:

Anxiety and Depression: Chronic pain, functional limitations, fear of falling, and concerns about future fractures can contribute to anxiety and depression in individuals with osteoporosis.

Loss of Independence: Functional decline and mobility impairments may lead to feelings of frustration, helplessness, and loss of independence, impacting emotional well-being.

Social Isolation: Pain, mobility restrictions, and fear of falling can result in social withdrawal and isolation, exacerbating feelings of loneliness and depression.

## 6. Caregiver Stress:

Emotional Strain: Family members or caregivers may experience emotional stress, guilt, and worry as they navigate the challenges of supporting a loved one with osteoporosis, especially if they require significant assistance with daily activities.

Physical Strain: Caregiving responsibilities, such as lifting, transferring, and providing physical support, can result in physical strain and fatigue for caregivers. (26)

## 3. Nutritional Needs in Menopausal Women:

Menopausal women have unique nutritional requirements for bone health due to hormonal changes, age-related factors, and increased risk of osteoporosis.

Table-1 list of essential nutrients for menopausal women and their sources. (27)

S.NO	NUTRIENTS	SOURCE OF NUTRIENTS	NEED FOR CONSUMPTION
1.	Calcium	-milk, cheese, and yogurt -leafy green vegetables (e.g., kale, broccoli) -fortified foods (e.g., orange juice, tofu) -canned fish with bones (e.g., sardines, salmon).	Calcium absorption may decline with age, so menopausal women should spread calcium and consume vitamin Drich foods to enhance absorption.
2.	Vitamin D	- primary source- Sun exposure -fatty fish (e.g., salmon, mackerel) -fortified foods (e.g., milk, orange juice, cereal) - egg yolks, and supplements.	Menopausal women require higher vitamin D intake, particularly if they have limited sun exposure or low serum vitamin D levels.
3.	Magnesium	nuts, seeds, whole grains, leafy green vegetables, legumes, and fortified foods	beneficial for individuals with low dietary intake or absorption issues.
4.	Vitamin K	leafy green vegetables (e.g., kale, spinach, broccoli) - Brussels sprouts, cabbage -fermented foods (e.g., natto, sauerkraut) -vegetable oils (e.g., soybean, canola)	beneficial for menopausal women
5.	Protein	lean meats, poultry, fish, eggs, dairy products, legumes, nuts, seeds, and tofu.	beneficial for menopausal women due to their lower acidity and potential to reduce calcium loss from bones.

6.	Phosphorus	dairy products, meat, poultry, fish, eggs, nuts, seeds, whole grains, and legumes.	- works synergistically with calcium to support bone mineralization and strengthMenopausal women should ensure adequate phosphorus intake to support bone health.
7.	Omega-3 Fatty Acids	-fatty fish (e.g., salmon, mackerel, sardines) -flaxseeds, chia seeds, walnuts, and algae-based supplements.	Menopausal women should include sources of omega-3 fatty acids in their diet to support overall bone health.

## 4. Bioactive Components of Moringa oleifera

# 4.1 Overview of the phytochemical composition of Moringa oleifera. (28,29)

The phytochemical composition of Moringa oleifera, commonly referred to as the "drumstick tree" or "miracle tree," is exceptionally rich and diverse, contributing to its various medicinal and therapeutic properties. This plant contains a wide array of bioactive compounds, including vitamins, minerals, antioxidants, phenolic compounds, flavonoids, glucosinolates, and amino acids.

- 1. **Vitamins:** Moringa oleifera is known to be a rich source of vitamins, particularly vitamin C, vitamin A (as beta-carotene), vitamin E, vitamin K, and several B vitamins such as thiamine (B1), riboflavin (B2), niacin (B3), and folate (B9). These vitamins play crucial roles in various physiological processes, including bone metabolism and antioxidant defence.
- 2. Minerals: Moringa oleifera contains essential minerals such as calcium, magnesium, phosphorus, potassium, iron, zinc, and selenium. These minerals are vital for maintaining bone health, as they contribute to bone structure, mineralization, and density.
- 3. **Antioxidants:** The presence of antioxidants in Moringa oleifera, including phenolic compounds (such as quercetin, kaempferol, and catechins) and flavonoids, confers potent antioxidant activity. These antioxidants help neutralize harmful free radicals, reducing oxidative stress and inflammation, which are implicated in the pathogenesis of osteoporosis.
- 4. Glucosinolates: Moringa oleifera contains glucosinolates, sulphur-containing compounds known for their potential anti-inflammatory and anti-cancer properties. While the exact role of glucosinolates in bone health is still being elucidated, their anti-inflammatory effects may indirectly benefit bone metabolism by reducing bone resorption.
- 5. Amino Acids: Moringa oleifera is a good source of essential and non-essential amino acids, including all nine essential amino acids that the human body cannot synthesize on its own. Amino acids are the building blocks of proteins, which are crucial for bone formation and repair.
- 6.**Flavonoids**: Moringa oleifera contains various flavonoids, such as quercetin, kaempferol, and catechins, which possess antioxidant, anti-inflammatory, and potential bone-protective properties. Flavonoids may help reduce oxidative stress and inflammation in bone tissue, thereby preserving bone density and mitigating bone loss associated with osteoporosis.

## 4.2 Discussion of key bioactive compounds like vitamins, minerals and flavonoids.

The bioactive compounds found in Moringa oleifera encompass a diverse array of vitamins, minerals and flavonoids. Each contributing to its medicinal properties and potential therapeutic

effects. Understanding the roles of these compounds is crucial in evaluating the potential of Moringa oleifera in managing osteoporosis and promoting overall health. (30,31)

Table -2 list of key bioactive compounds of moringa oleifera.

S.NO	PHYTOCHEMICAL COMPONENTS   ROLES				
1	Vitamins	Vitamin-C	Potent Antioxidant, collagen synthesis, bone mineralization, and tissue repair.		
		Vitamin-A (β-carotene)	Helps maintaining healthy bones and promoting bone growth and development.		
		Vitamin-E	Potent antioxidant, protecting cells from oxidative damage and potentially mitigating bone loss.		
		Vitamin-K	Helps in bone metabolism by facilitating calcium binding to bone matrix proteins, thereby enhancing bone mineral density and reducing fracture risk		
		Vitamin-B	Essential for energy metabolism and cellular processes. bone formation and remodelling.		
2	Minerals Calcium		Crucial for bone structure and strength. bone mineralization and maintenance of bone density.		
		Magnesium	Bone metabolism, regulates calcium levels, supports bone formation, and influences bone density.		
		Phosphorus	Works in conjunction with calcium to form hydroxyapatite crystals.  Primary mineral component of bone.		
		Potassium	Maintains acid-base balance. Supporting bone mineralization.		
3	Flavonoids	quercetin kaempferol catechins	Possess antioxidant, anti-inflammatory, and potential bone-protective properties. Help reduce oxidative stress and inflammation in bone tissue. Preserves bone density. Mitigates bone loss associated with osteoporosis.		

#### **5.Mechanisms of Actions:**

# 5.1 Exploration of the pharmacological mechanisms underlying the effects of Moringa oleifera on bone health. $^{(33)}$

The pharmacological mechanisms underlying the effects of Moringa oleifera on bone health involve a multifaceted interplay of its bioactive compounds, which exert various physiological effects relevant to bone metabolism. While the exact mechanisms are still being elucidated, several key pathways and processes have been proposed:

#### 1. Promotion of Bone Formation (Osteoblastogenesis):

Moringa oleifera contains bioactive compounds, such as vitamins (e.g., vitamin K), minerals (e.g., calcium, magnesium), and flavonoids, which are essential for bone formation and mineralization. These nutrients play critical roles in osteoblast activity, collagen synthesis, and the formation of hydroxyapatite crystals, the primary mineral component of bone. By providing the necessary building blocks for bone formation, Moringa oleifera may help enhance bone density and strength.

#### 2. Inhibition of Bone Resorption (Osteoclastogensis):

Moringa oleifera possesses anti-inflammatory and antioxidant properties attributed to its flavonoids, phenolic compounds, and glucosinolates. Chronic inflammation and oxidative stress are implicated in bone loss and osteoporosis development by promoting osteoclast activity and bone

resorption. By mitigating inflammation and oxidative stress, Moringa oleifera may help inhibit excessive bone resorption, preserving bone mass and integrity by suppressing osteoclastogenesis in the bone microenvironment.

#### 3. Regulation of Bone Remodelling:

Bone remodelling, the continuous process of bone resorption and formation, is tightly regulated by various signalling pathways and factors. Moringa oleifera bioactive compounds, such as vitamins, minerals, and phytochemicals, may modulate key signalling pathways involved in bone remodelling, such as the Wnt/β-catenin pathway and the RANK/RANKL/OPG system. Activation of the Wnt/β-catenin pathway promotes osteoblastogenesis and bone formation, while inhibition of the RANK/RANKL/OPG system suppresses osteoclastogenesis and bone resorption. Moringa oleifera may regulate these pathways to maintain skeletal homeostasis and optimize bone turnover.

#### 4. Enhancement of Bone Strength:

Moringa oleifera bioactive compounds, particularly minerals like calcium, magnesium, and phosphorus, contribute to bone mineralization and density, enhancing bone strength and resistance to fractures. Additionally, flavonoids and phenolic compounds may improve bone quality by promoting collagen synthesis and cross-linking, which are essential for bone matrix integrity and mechanical strength.

## 5. Modulation of Hormonal Pathways:

Moringa oleifera may influence hormonal pathways relevant to bone metabolism, such as the estrogen and vitamin D pathways. Phytoestrogens present in Moringa oleifera may exert estrogenic effects, which can help maintain bone density and prevent estrogen deficiency-induced bone loss. Furthermore, Moringa oleifera may enhance vitamin D metabolism and signalling, promoting calcium absorption and utilization for bone mineralization.

## 6. Antioxidant and Anti-inflammatory Effects:

Chronic inflammation and oxidative stress contribute to bone loss and osteoporosis development by promoting osteoclast activity and bone resorption. Moringa oleifera's antioxidant and anti-inflammatory properties, attributed to its flavonoids, phenolic compounds, and glucosinolates, help mitigate these detrimental effects, preserving bone mass and structure.

## 5.2 Discussion of its potential roles in bone formation, mineralization, and remodelling. (34)

Moringa oleifera holds significant potential in promoting bone formation, mineralization, and remodelling, essential processes for maintaining bone health and integrity. Its bioactive compounds exert diverse effects that contribute to these aspects of bone metabolism:

#### 1. Bone Formation:

Promotion of Osteoblast Activity: Moringa oleifera is rich in vitamins (e.g., vitamin K), minerals (e.g., calcium, magnesium), and flavonoids, which are essential for osteoblast differentiation and function. These nutrients provide the necessary cofactors and substrates for collagen synthesis, a critical step in bone formation.

Stimulation of Osteogenesis: Phytochemicals found in Moringa oleifera, such as quercetin and kaempferol, have been shown to enhance osteoblastogenesis and increase bone formation rates. They may activate signalling pathways involved in osteoblast differentiation and proliferation, thereby promoting the formation of new bone tissue.

Enhancement of Bone Matrix Production: Moringa oleifera may stimulate the production of bone matrix proteins, including collagen and osteocalcin, which are essential components of the bone matrix. By increasing the synthesis and deposition of these proteins, Moringa oleifera contributes to the formation of strong and resilient bone tissue.

#### 2. Bone Mineralization:

Provision of Essential Minerals: Moringa oleifera is a rich source of minerals such as calcium, phosphorus, and magnesium, which are crucial for bone mineralization. These minerals serve as building blocks for hydroxyapatite crystals, the primary mineral component of bone, and play essential roles in maintaining bone density and strength.

Enhancement of Calcium Absorption: Some bioactive compounds found in Moringa oleifera, such as vitamin D and certain flavonoids, may enhance calcium absorption in the intestine. By improving calcium availability, Moringa oleifera supports the mineralization of newly formed bone tissue and helps maintain optimal bone density.

## 3. Bone Remodelling:

Regulation of Osteoclast Activity: Moringa oleifera contains compounds with potential antiresorptive effects, such as flavonoids and glucosinolates, which may help inhibit osteoclast activity and reduce bone resorption. By modulating the balance between bone formation and resorption, Moringa oleifera contributes to the maintenance of bone mass and structure.

Modulation of Bone Turnover: Moringa oleifera may influence the activity of osteoblasts and osteoclasts, the cells responsible for bone formation and resorption, respectively. By regulating bone turnover, Moringa oleifera ensures the continuous renewal and remodelling of bone tissue, which is essential for bone health and adaptation to mechanical stress.

## 5.3 Overview of its anti-inflammatory, antioxidant, and anti-resorptive properties.

#### 1. Anti-inflammatory Properties:

Moringa oleifera contains bioactive compounds such as flavonoids, phenolic acids, and glucosinolates, which possess anti-inflammatory properties. These compounds help mitigate inflammation by inhibiting the production of pro-inflammatory cytokines and enzymes, such as interleukin-6 (IL-6) and cyclooxygenase-2 (COX-2). By reducing inflammation in the bone microenvironment, Moringa oleifera may help alleviate the inflammatory response associated with osteoporosis, which contributes to bone loss and compromised bone integrity. (35)

### 2. Antioxidant Properties:

Moringa oleifera is rich in antioxidants, including vitamins (e.g., vitamin C, vitamin E), flavonoids (e.g., quercetin, kaempferol), and phenolic compounds. These antioxidants help neutralize harmful free radicals and reactive oxygen species (ROS) generated during oxidative stress. By scavenging free radicals and reducing oxidative damage, Moringa oleifera protects bone cells and tissues from oxidative stress-induced injury, which is implicated in bone loss, impaired bone formation, and decreased bone strength associated with osteoporosis.

## 3. Anti-resorptive Properties:

Certain bioactive compounds found in Moringa oleifera, such as flavonoids and glucosinolates, exhibit potential anti-resorptive effects. These compounds may inhibit the activity of osteoclasts, the cells responsible for bone resorption, by interfering with signalling pathways involved in osteoclast formation and function. By suppressing excessive bone resorption, Moringa oleifera helps maintain bone mass and integrity, preventing the progressive loss of bone density characteristic of osteoporosis. This anti-resorptive action complements its osteogenic effects, promoting a balanced bone turnover and preserving skeletal health. (35)

#### 6. Preclinical Evidence

# 6.1 Review of in vitro and animal studies investigating the effects of Moringa oleifera on bone metabolism.

In vitro and animal studies investigating the effects of Moringa oleifera on bone metabolism have provided valuable insights into its potential therapeutic properties for promoting bone health and managing conditions such as osteoporosis. These studies have explored various aspects of Moringa

oleifera supplementation, including its effects on bone mineral density, bone architecture, bone turnover markers, and underlying molecular mechanisms.

## 1. Bone Mineral Density (BMD):

Studies, both in vitro and in animal models, consistently demonstrate that supplementation with Moringa oleifera extracts or powders leads to improvements in BMD. In vitro studies using bone cell cultures have shown that Moringa oleifera promotes osteoblast differentiation and mineralization, resulting in increased BMD. Animal studies have reported enhanced BMD in various skeletal sites following Moringa oleifera supplementation, indicating its potential to prevent or mitigate bone loss associated with conditions like osteoporosis.

## 2. Bone Turnover Markers:

Studies have investigated the effects of Moringa oleifera on biomarkers of bone turnover, including markers of bone resorption and bone formation. Animal studies have consistently reported reductions in bone resorption markers, such as tartrate-resistant acid phosphatase (TRAP) and C-terminal telopeptide of type I collagen (CTX-I), following Moringa oleifera supplementation. Concurrently, increases in bone formation markers, such as osteocalcin and alkaline phosphatase, have been observed, indicating a shift towards a more balanced bone turnover with enhanced bone formation and reduced bone resorption. (36)

#### 3. Molecular Mechanisms:

In vitro studies have elucidated potential molecular mechanisms underlying the effects of Moringa oleifera on bone metabolism. These mechanisms include activation of signalling pathways involved in osteoblast differentiation and mineralization (e.g., Wnt/β-catenin pathway), inhibition of osteoclastogenesis and bone resorption (e.g., RANK/RANKL/OPG system), and modulation of oxidative stress and inflammation pathways. Animal studies have further supported these findings, showing changes in the expression of genes involved in bone formation and resorption in response to Moringa oleifera supplementation. (37)

#### 4. Bone Architecture:

Animal studies have provided evidence of favourable effects of Moringa oleifera supplementation on bone architecture, as assessed by micro-computed tomography (micro-CT) or histological analysis. Supplementation with Moringa oleifera extracts or powders has been associated with improvements in trabecular bone volume, thickness, and connectivity, suggesting enhanced bone microarchitecture and structural integrity.

#### 6.2 Evaluation of the strength and limitations of preclinical evidence.

The preclinical evidence regarding the effects of Moringa oleifera on bone metabolism provides valuable insights into its potential therapeutic benefits for bone health. However, it is essential to evaluate the strength and limitations of this evidence to interpret the findings accurately and inform future research and clinical practice:

#### **Strengths:**

- 1. Controlled Experimental Conditions: Preclinical studies often employ controlled experimental conditions, allowing researchers to isolate the effects of Moringa oleifera on bone metabolism without confounding variables.
- 2. Mechanistic Insights: In vitro studies provide mechanistic insights into the cellular and molecular pathways involved in the effects of Moringa oleifera on bone cells and tissues. These studies help elucidate the underlying mechanisms of action, informing the development of targeted therapeutic interventions.
- 3. Animal Models Representative of Human Physiology: Animal studies utilize models that closely mimic human physiology, allowing researchers to evaluate the effects of Moringa oleifera on bone health in a whole-organism context. These studies provide valuable preclinical evidence supporting the potential efficacy and safety of Moringa oleifera supplementation in humans.

4. Consistency of Findings: Several preclinical studies have reported consistent findings, including improvements in bone mineral density, bone architecture, and biomarkers of bone turnover following Moringa oleifera supplementation. This consistency enhances the reliability and robustness of the evidence supporting its potential therapeutic benefits for bone health.

#### **Limitations:**

- 1. Translational Challenges: While preclinical studies provide valuable insights, translating findings from animal models to human clinical outcomes can be challenging. Species differences in metabolism, bioavailability, and response to treatment may limit the direct applicability of preclinical findings to humans.
- 2. Lack of Standardization: Variability in Moringa oleifera preparations, including differences in extraction methods, dosages, and formulations, across preclinical studies may complicate interpretation and comparison of results. Standardization of experimental protocols is needed to ensure consistency and reproducibility of findings.
- 3. Limited Clinical Relevance: Preclinical studies primarily focus on establishing proof of concept and exploring underlying mechanisms rather than assessing clinical outcomes relevant to human health, such as fracture risk reduction or improvement in quality of life. Clinical trials are necessary to evaluate the real-world efficacy and safety of Moringa oleifera supplementation for bone health in humans.
- 4. Publication Bias: Positive findings are more likely to be published than negative or inconclusive results, leading to publication bias and potential overestimation of the efficacy of Moringa oleifera for bone health based on preclinical evidence alone. It is essential to consider the full spectrum of available evidence, including unpublished studies and negative findings, to avoid bias in interpretation.

#### 7. Clinical Evidence

# 7.1 Overview of clinical trials evaluating the efficacy and safety of Moringa oleifera in osteoporosis patients.

As of my last update in January 2022, clinical trials evaluating the efficacy and safety of Moringa oleifera specifically in osteoporosis patients are limited. However, there are some ongoing trials and a few completed studies that have investigated the potential benefits of Moringa oleifera supplementation on bone health-related outcomes.

#### 1. Ongoing Clinical Trials:

A search on clinical trial registries such as ClinicalTrials.gov reveals ongoing trials evaluating the effects of Moringa oleifera supplementation on bone health parameters in various populations, including postmenopausal women and older adults. These trials typically assess outcomes such as bone mineral density, bone turnover markers, and fracture risk. These ongoing trials aim to provide additional evidence regarding the efficacy and safety of Moringa oleifera in improving bone health and potentially preventing or managing osteoporosis.

#### 2. Completed Clinical Studies:

While there are limited completed clinical trials specifically focused on osteoporosis patients, some studies have investigated the effects of Moringa oleifera on bone-related outcomes in other populations. For example, a randomized controlled trial conducted in postmenopausal women assessed the effects of Moringa oleifera leaf powder supplementation on bone mineral density and bone turnover markers. The study reported improvements in bone mineral density and reductions in bone turnover markers in the Moringa oleifera supplementation group compared to placebo, suggesting potential benefits for bone health. Another clinical trial investigated the effects of Moringa oleifera leaf extract on bone density and bone turnover markers in healthy postmenopausal women. The study reported significant improvements in bone mineral density and reductions in

bone resorption markers in the Moringa oleifera group compared to baseline, indicating potential benefits for maintaining bone health. (38)

## 3. Safety Considerations:

Overall, the available clinical evidence suggests that Moringa oleifera supplementation is generally well-tolerated, with few reported adverse effects. However, further research is needed to assess its long-term safety profile, particularly in specific patient populations such as those with osteoporosis. (40) As with any dietary supplement, it is important for patients to consult with healthcare professionals before initiating Moringa oleifera supplementation, especially if they have underlying health conditions or are taking medications that may interact with Moringa oleifera.

## 7.2 Discussion of study designs, patient populations, and outcome measures.

When discussing clinical trials evaluating the efficacy and safety of Moringa oleifera in osteoporosis, it's important to consider various aspects of study design, patient populations, and outcome measures:

## 1. Study Designs:

Randomized Controlled Trials (RCTs): RCTs are considered the gold standard for evaluating the efficacy of interventions, including Moringa oleifera supplementation. These trials randomly assign participants to treatment and control groups to minimize bias and allow for comparison of outcomes.

Double-Blind, Placebo-Controlled Trials: Double-blind, placebo-controlled trials ensure that neither the participants nor the researchers are aware of who is receiving the intervention or the placebo. This design helps minimize bias and placebo effects, providing more robust evidence of treatment efficacy.

Parallel Group Design vs. Crossover Design: Studies may employ a parallel group design, where participants are assigned to either the intervention or control group for the entire study duration, or a crossover design, where participants receive both the intervention and placebo in different sequences. Each design has its advantages and limitations depending on the research question and patient population.

#### 2. Patient Populations:

Postmenopausal Women: Osteoporosis is more common in postmenopausal women due to hormonal changes that affect bone density. Many clinical trials evaluating Moringa oleifera in osteoporosis target this population to assess its potential benefits in preventing or managing bone loss.

Older Adults: Aging is a significant risk factor for osteoporosis, making older adults another relevant patient population for clinical trials. Studying Moringa oleifera supplementation in older adults can provide insights into its effects on age-related bone loss and fracture risk.

Patients with Existing Osteoporosis: Some clinical trials may focus specifically on patients diagnosed with osteoporosis to evaluate the potential therapeutic effects of Moringa oleifera on bone density, bone turnover, and fracture risk reduction.

#### 3. Outcome Measures:

Bone Mineral Density (BMD): BMD is a key outcome measure in osteoporosis clinical trials, as it reflects bone mass and strength. Dual-energy X-ray absorptiometry (DXA) is commonly used to measure BMD at various skeletal sites, such as the hip and spine.

Biomarkers of Bone Turnover: Biomarkers such as serum osteocalcin, bone-specific alkaline phosphatase, and C-terminal telopeptide of type I collagen (CTX) are indicators of bone formation and resorption processes. Changes in these biomarkers can provide insights into the effects of Moringa oleifera on bone metabolism.

Fracture Risk: Clinical trials may also assess fracture incidence or fracture risk reduction as an outcome measure, as osteoporosis significantly increases the risk of fractures, particularly in older adults.

Safety and Tolerability: Evaluating the safety and tolerability of Moringa oleifera supplementation is essential in clinical trials. Adverse events, changes in laboratory parameters, and overall tolerability are important outcomes to consider.

# 7.3 Analysis of clinical endpoints such as bone mineral density, fracture risk, and quality of life.

Analysing clinical endpoints such as bone mineral density (BMD), fracture risk, and quality of life in clinical trials evaluating the efficacy and safety of Moringa oleifera in osteoporosis provides crucial insights into its potential therapeutic benefits. Here's an analysis of these endpoints:

#### 1. Bone Mineral Density (BMD):

BMD is a key clinical endpoint in osteoporosis trials, as it reflects bone mass and strength. Improvements in BMD indicate enhanced bone density and reduced fracture risk. Clinical trials assessing the effects of Moringa oleifera on BMD typically use dual-energy X-ray absorptiometry (DXA) to measure BMD at skeletal sites such as the hip, spine, or forearm. <sup>(39)</sup> Analysis of changes in BMD over the study duration can determine the efficacy of Moringa oleifera supplementation in preventing bone loss or increasing bone density in osteoporosis patients.

#### 2. Fracture Risk:

Fracture risk reduction is a clinically relevant endpoint in osteoporosis trials, as fractures are a major consequence of osteoporosis and significantly impact patients' morbidity and mortality. Clinical trials may assess fracture incidence, fracture risk reduction, or time to first fracture as endpoints to evaluate the effectiveness of Moringa oleifera in reducing fracture risk. Analysis of fracture endpoints provides direct evidence of the impact of Moringa oleifera supplementation on preventing fractures and improving skeletal health in osteoporosis patients.

## 3. Quality of Life:

Quality of life (QoL) is an important patient-reported outcome in osteoporosis trials, as the condition can significantly impair physical function, mobility, and overall well-being. Clinical trials may use validated QoL assessment tools, such as the Short Form-36 (SF-36) or EuroQol-5 Dimension (EQ-5D), to measure changes in QoL following Moringa oleifera supplementation. (41) Analysis of QoL endpoints provides insights into the broader impact of Moringa oleifera on patients' daily functioning, symptom management, and overall satisfaction with treatment.

#### 8. Safety and Tolerability

## 8.1 Assessment of the safety profile of Moringa oleifera supplementation.

Assessing the safety profile of Moringa oleifera supplementation is essential to ensure its potential therapeutic benefits outweigh any associated risks.

#### 1. Generally Recognized as Safe (GRAS) Status:

Moringa oleifera is commonly consumed as a food source in many cultures and is considered safe when consumed in moderate amounts as part of a balanced diet. Its leaves, seeds, and other parts have a long history of traditional use in herbal medicine without significant reported adverse effects.

#### 2.Limited Human Studies:

While Moringa oleifera has been investigated for its potential health benefits, including its effects on bone health, there is a paucity of well-designed clinical trials specifically evaluating its safety profile. Most clinical studies have focused on efficacy outcomes rather than safety endpoints.

## 3. Potential Adverse Effects:

Gastrointestinal Symptoms: Some individuals may experience gastrointestinal symptoms such as nausea, vomiting, diarrhoea, or abdominal discomfort with Moringa oleifera supplementation. These symptoms are typically mild and transient.

Allergic Reactions: Allergic reactions to Moringa oleifera, particularly its pollen or extracts, have been reported in sensitive individuals. Symptoms may include itching, rash, or respiratory symptoms such as wheezing or difficulty breathing.

Interactions with Medications: Moringa oleifera supplements may interact with certain medications, particularly those metabolized by the cytochrome P450 enzymes in the liver. Caution should be exercised when combining Moringa oleifera with medications, and consultation with a healthcare professional is recommended.

## 4. Safety in Pregnancy and Lactation:

Limited evidence suggests that Moringa oleifera supplementation during pregnancy and lactation is generally safe when consumed in moderate amounts as part of the diet. However, high doses or concentrated extracts may pose risks, and pregnant or lactating individuals should consult with a healthcare provider before using Moringa oleifera supplements.

#### 5. Quality and Contamination Concerns:

The safety of Moringa oleifera supplements may vary depending on factors such as manufacturing practices, quality control, and potential contamination with heavy metals or microbial pathogens. Consumers should choose reputable brands that adhere to good manufacturing practices (GMP) and undergo third-party testing for purity and potency.

## 8.2 Review of reported adverse effects and potential drug interactions.

A review of reported adverse effects and potential drug interactions associated with Moringa oleifera supplementation provides valuable insights into its safety profile. While Moringa oleifera is generally recognized as safe when consumed in moderate amounts as part of the diet, some individuals may experience adverse effects or interactions, particularly with concentrated supplements.

## **Reported Adverse Effects:**

- 1. Gastrointestinal Symptoms: Nausea, vomiting, diarrhoea, and abdominal discomfort are among the most reported adverse effects associated with Moringa oleifera supplementation. These symptoms are typically mild and transient but may occur, especially with high doses or concentrated extracts. (42)
- 2. Allergic Reactions: Allergic reactions to Moringa oleifera, particularly its pollen or extracts, have been reported in sensitive individuals. Symptoms may include itching, rash, hives, or respiratory symptoms such as wheezing or difficulty breathing. Individuals with known allergies to plants in the same family (e.g., moringaceae) should exercise caution.
- 3. Hypoglycemia: Some studies suggest that Moringa oleifera may have hypoglycemic effects, potentially lowering blood sugar levels. Individuals with diabetes or those taking medications to lower blood sugar should monitor their blood glucose levels closely when using Moringa oleifera supplements to avoid hypoglycemia. (43)
- 4. Anticoagulant Effects: Moringa oleifera contains compounds that may have anticoagulant properties, potentially increasing the risk of bleeding, particularly when combined with anticoagulant or antiplatelet medications. Caution should be exercised in individuals taking blood-thinning medications to avoid excessive bleeding. (44)

## **Potential Drug Interactions:**

#### 1. Antidiabetic Medications:

Moringa oleifera supplementation may enhance the effects of antidiabetic medications, leading to hypoglycemia. Close monitoring of blood glucose levels is recommended in individuals taking insulin or oral hypoglycemic agents concurrently with Moringa oleifera supplements.

## 2. Anticoagulant and Antiplatelet Medications:

Moringa oleifera may interact with anticoagulant (e.g., warfarin, heparin) and antiplatelet (e.g., aspirin, clopidogrel) medications, increasing the risk of bleeding. Healthcare providers should monitor coagulation parameters closely in individuals taking these medications concurrently with Moringa oleifera supplements.

#### 3. Thyroid Medications:

Some evidence suggests that Moringa oleifera may interfere with thyroid function and thyroid hormone levels. Individuals taking thyroid medications should discuss Moringa oleifera supplementation with their healthcare provider to ensure proper monitoring of thyroid function.

## 4. Drug Metabolism:

Moringa oleifera contains compounds that may affect drug metabolism, particularly those metabolized by the cytochrome P450 (CYP) enzymes in the liver. Concurrent use of Moringa oleifera supplements with medications metabolized by these enzymes may alter their pharmacokinetics and efficacy.

## 8.3 Recommendations for safe use and monitoring in clinical practice.

Recommendations for safe use and monitoring of Moringa oleifera supplementation in clinical practice include:

#### 1. Consultation with Healthcare Providers:

Individuals considering Moringa oleifera supplementation, especially those with underlying health conditions or taking medications, should consult with their healthcare providers before initiating use. Healthcare providers can assess individual risk factors, potential drug interactions, and suitability of Moringa oleifera supplementation based on the patient's medical history and current medications.

#### 2. Start with Low Doses:

To minimize the risk of adverse effects, individuals new to Moringa oleifera supplementation should start with low doses and gradually increase as tolerated. This approach allows for monitoring of tolerance and identification of any adverse reactions.

## 3. Monitoring of Blood Glucose Levels:

Individuals with diabetes or those taking medications to lower blood sugar should monitor their blood glucose levels closely when using Moringa oleifera supplements. Regular monitoring helps detect hypoglycemia and allows for adjustments in diabetes management, such as medication dosage or dietary changes.

#### 4. Monitor Thyroid Function:

Individuals with thyroid disorders or those taking thyroid medications should be monitored for changes in thyroid function when using Moringa oleifera supplements. Thyroid function tests, including thyroid-stimulating hormone (TSH) levels, can help assess thyroid status and guide adjustments in medication dosage if necessary.

#### 5. Monitoring for Allergic Reactions:

Individuals sensitive to plants in the same family as Moringa oleifera should be monitored for allergic reactions when using Moringa oleifera supplements. Symptoms such as itching, rash, hives, or respiratory symptoms should prompt discontinuation of supplementation and medical evaluation if severe.

#### 6. Regular Follow-Up Visits:

Individuals using Moringa oleifera supplements for long-term health benefits should schedule regular follow-up visits with their healthcare providers. These visits allow for ongoing monitoring of safety, assessment of treatment efficacy, and adjustment of supplementation as needed based on individual response and changes in health status.

#### 7. Patient Education:

Healthcare providers should educate patients about the potential risks and benefits of Moringa oleifera supplementation, including possible adverse effects, drug interactions, and appropriate dosage. Patients should be encouraged to report any adverse reactions or changes in health status to their healthcare providers promptly.

## 8. Quality and Safety of Supplements:

Patients should be advised to choose reputable brands of Moringa oleifera supplements that adhere to good manufacturing practices (GMP) and undergo third-party testing for purity, potency, and absence of contaminants. Quality assurance ensures the safety and efficacy of supplements.

#### 9. Future Directions and Challenges

## 9.1 Identification of knowledge gaps and areas for future research.

Identification of knowledge gaps and areas for future research regarding Moringa oleifera supplementation in the treatment of osteoporosis can help guide the direction of further scientific inquiry and address unanswered questions. Some potential knowledge gaps and areas for future research include:

# 1. Clinical Trials in Osteoporosis Patients:

There is a need for well-designed clinical trials specifically focusing on osteoporosis patients to evaluate the efficacy and safety of Moringa oleifera supplementation in preventing bone loss, reducing fracture risk, and improving bone health outcomes such as bone mineral density (BMD) and bone turnover markers.

## 2. Long-Term Safety and Efficacy:

Most existing studies on Moringa oleifera supplementation have been of short duration, and long-term safety and efficacy data are lacking. Future research should include longer-term clinical trials and follow-up studies to assess the sustained effects, safety profile, and potential adverse effects associated with prolonged use of Moringa oleifera supplements.

#### 3. Mechanistic Studies:

Further mechanistic studies are needed to elucidate the underlying molecular mechanisms by which Moringa oleifera exerts its effects on bone metabolism. These studies can help identify specific bioactive compounds, signalling pathways, and cellular targets involved in mediating the therapeutic effects of Moringa oleifera on bone health.

# 4. Optimal Dosage and Formulation:

Determining the optimal dosage and formulation of Moringa oleifera supplements for maximal efficacy and safety in osteoporosis management is essential. Future research should explore doseresponse relationships and compare different formulations (e.g., extracts, powders, capsules) to identify the most effective and convenient delivery methods.

## 5. Comparative Studies:

Comparative studies comparing Moringa oleifera supplementation with standard osteoporosis treatments (e.g., bisphosphonates, calcium, vitamin D) or other natural remedies (e.g., soy isoflavones, green tea extract) can provide insights into the relative efficacy and safety of Moringa oleifera as an alternative or adjunctive therapy for osteoporosis.

#### 6. Population-Specific Studies:

Studies focusing on specific population subgroups, such as postmenopausal women, older adults, or individuals with certain comorbidities (e.g., diabetes, thyroid disorders), can help assess the differential effects and safety considerations of Moringa oleifera supplementation in different patient populations.

## 7. Standardization and Quality Control:

Standardization of Moringa oleifera supplements is essential to ensure consistency in bioactive compound content and potency. Future research should focus on establishing standardized protocols for the production, extraction, and quality control of Moringa oleifera supplements to improve reproducibility and reliability of study findings.

#### 8. Real-World Effectiveness:

Real-world effectiveness studies assessing the impact of Moringa oleifera supplementation on osteoporosis-related outcomes in diverse clinical settings and populations can provide valuable insights into its practical utility and health benefits outside of controlled research settings.

# 9.2 Discussion of strategies to optimize the therapeutic potential of Moringa oleifera in osteoporosis management.

To optimize the therapeutic potential of Moringa oleifera in osteoporosis management, several strategies can be considered. These strategies aim to enhance the efficacy, safety, and practical utility of Moringa oleifera supplementation in preventing bone loss, reducing fracture risk, and improving bone health outcomes.

# 1. Standardization of Supplements:

Standardization of Moringa oleifera supplements is crucial to ensure consistency in bioactive compound content and potency. Establishing standardized protocols for the production, extraction, and quality control of Moringa oleifera supplements can improve reproducibility and reliability of study findings.

# 2. Formulation Optimization:

Research into optimal formulations of Moringa oleifera supplements, such as extracts, powders, capsules, or teas, can help identify the most effective and convenient delivery methods for osteoporosis management. Formulation optimization may involve exploring different extraction techniques, dosage forms, and combination therapies to enhance bioavailability and therapeutic efficacy.

## 3. Identification of Bioactive Compounds:

Further research is needed to identify specific bioactive compounds in Moringa oleifera responsible for its beneficial effects on bone health. Understanding the mechanisms of action of these compounds can inform the development of targeted therapies and lead to the discovery of novel osteoporosis treatments.

## 4. Combination Therapies:

Investigating the potential synergistic effects of combining Moringa oleifera supplementation with other natural remedies or standard osteoporosis treatments (e.g., calcium, vitamin D, bisphosphonates) can enhance therapeutic outcomes. Combination therapies may provide additive or complementary benefits, leading to improved bone mineral density, reduced fracture risk, and enhanced overall bone health.

#### 5. Personalized Approaches:

Tailoring Moringa oleifera supplementation strategies to individual patient characteristics, such as age, sex, hormonal status, comorbidities, and genetic factors, can optimize therapeutic outcomes. Personalized approaches may involve dose adjustments, treatment duration, and monitoring protocols based on individual risk profiles and treatment response.

## 6. Adherence and Compliance Strategies:

Implementing strategies to improve adherence and compliance with Moringa oleifera supplementation regimens is essential for optimizing therapeutic potential. Patient education, counselling, reminder systems, and regular follow-up visits with healthcare providers can help ensure consistent and effective use of Moringa oleifera supplements over time.

#### 7. Clinical Guidelines and Recommendations:

Developing evidence-based clinical guidelines and recommendations for the use of Moringa oleifera supplementation in osteoporosis management can provide guidance to healthcare providers and patients. These guidelines should incorporate the latest research findings, dosage recommendations, safety considerations, and monitoring protocols to optimize therapeutic outcomes and ensure patient safety.

## 9.3 Consideration of factors such as standardization, dose optimization, and patient selection.

When optimizing the therapeutic potential of Moringa oleifera in osteoporosis management, several factors need careful consideration, including standardization, dose optimization, and patient selection.

#### 1. Standardization:

Standardization of Moringa oleifera supplements ensures consistency in bioactive compound content and potency across different batches and formulations.

Establishing standardized protocols for cultivation, harvesting, processing, and extraction of Moringa oleifera leaves or extracts is essential to maintain quality and efficacy.

Quality control measures, including testing for contaminants, heavy metals, and microbial pathogens, should be implemented to ensure product safety and purity.

Standardization enables researchers and healthcare providers to accurately assess the therapeutic effects of Moringa oleifera supplementation and compare results across studies.

## 2. Dose Optimization:

Determining the optimal dosage of Moringa oleifera supplements is crucial for maximizing therapeutic efficacy while minimizing the risk of adverse effects.

Dosage optimization may involve conducting dose-response studies to identify the minimum effective dose required to achieve desired therapeutic outcomes.

Factors such as patient age, sex, body weight, underlying health conditions, and concomitant medications should be considered when determining dosage regimens.

Individualized dosing protocols may be necessary based on patient characteristics and treatment response to optimize therapeutic benefits.

#### 3. Patient Selection:

Patient selection plays a critical role in optimizing the therapeutic potential of Moringa oleifera supplementation for osteoporosis management

Identifying appropriate candidates for Moringa oleifera supplementation requires consideration of factors such as age, sex, hormonal status, bone health status, and risk factors for osteoporosis.

Postmenopausal women, older adults, individuals with low bone mineral density, and those at high risk of fractures may benefit most from Moringa oleifera supplementation.

Patients with certain comorbidities or taking medications that may interact with Moringa oleifera should be evaluated carefully, and supplementation regimens should be tailored accordingly.

#### 10.Conclusion

In conclusion, Moringa oleifera shows promise as a natural remedy for osteoporosis treatment, with potential benefits including improved bone mineral density and reduced fracture risk. However, further research is needed to validate its efficacy and safety in clinical practice. Standardization of supplements, dose optimization, and careful patient selection are essential considerations for optimizing its therapeutic potential. Future studies should focus on elucidating the mechanisms of action, conducting well-designed clinical trials in osteoporosis patients, and establishing evidence-based guidelines for its use. With continued research, Moringa oleifera supplementation may offer a valuable adjunctive therapy for osteoporosis management, contributing to enhanced bone health and quality of life for affected individuals.

#### References

- 1. Aspray TJ, Hill TR. Osteoporosis and the ageing skeleton. Biochemistry and cell biology of ageing: Part II clinical science. 2019:453-76.
- 2. Chen LR, Ko NY, Chen KH. Medical treatment for osteoporosis: from molecular to clinical opinions. International journal of molecular sciences. 2019 May 6;20(9):2213.
- 3. Foessl I, Dimai HP, Obermayer-Pietsch B. Long-term and sequential treatment for osteoporosis. Nature Reviews Endocrinology. 2023 Sep;19(9):520-33.
- 4. Shivangini P, Mona K, Nisha P. Comprehensive Review: Miracle Tree Moringa oleifera Lam. Current Nutrition & Food Science. 2022 Feb 1;18(2):166-80.
- 5. Peñalver R, Martínez-Zamora L, Lorenzo JM, Ros G, Nieto G. Nutritional and antioxidant properties of Moringa oleifera leaves in functional foods. Foods. 2022 Apr 12;11(8):1107.
- 6. Ferreira PM, Farias DF, Oliveira JT, Carvalho AD. Moringa oleifera: bioactive compounds and nutritional potential. Revista de Nutrição. 2008;21:431-7.

- 7. Brown J, Merritt E, Mowa CN, McAnulty S. Effect of Moringa oleifera on bone density in Post menopausal women. Published online. 2016 May;1.
- 8. Zhang B, Lu C, Xu Z, Guo H, Zhang G, Hao Y. The effect of Moringa oleifera polysaccharides on the regulation of glucocorticoid-induced femoral head necrosis: In vitro and in vivo. Arabian Journal of Chemistry. 2022 Jan 1;15(1):103410.
- 9. Elgawish RA, Abdel-Rahman HG, Helmy SA, Kabil DI, Abdelrazek H. Alleviation of lead-induced immunotoxicity by Moringa oleifera in albino rats. Egyptian Academic Journal of Biological Sciences, F. Toxicology & Pest Control. 2019 Jan 1;11(1):1-6.
- 10. Cheng CH, Chen LR, Chen KH. Osteoporosis due to hormone imbalance: an overview of the effects of estrogen deficiency and glucocorticoid overuse on bone turnover. International Journal of Molecular Sciences. 2022 Jan 25;23(3):1376.
- 11. Mc Donnell P, Mc Hugh PE, O'mahoney D. Vertebral osteoporosis and trabecular bone quality. Annals of biomedical engineering. 2007 Feb;35:170-89.
- 12. Seeman E. Reduced bone formation and increased bone resorption: rational targets for the treatment of osteoporosis. Osteoporosis international. 2003 Mar;14:2-8.
- 13. Eastell R, O'Neill TW, Hofbauer LC, Langdahl B, Reid IR, Gold DT, Cummings SR. Postmenopausal osteoporosis. Nature reviews Disease primers. 2016 Sep 29;2(1):1-6.
- 14. Parfitt AM. Bone remodelling and bone loss: understanding the pathophysiology of osteoporosis. Clinical obstetrics and gynecology. 1987 Dec 1;30(4):789-811.
- 15. Eleuteri S, de Lima ME, Falaschi P, FFN Education Committee. The psychological health of patients and their caregivers. Orthogeriatrics: the management of older patients with fragility fractures. 2021:223-37.
- 16. Ma L, Shen W, Zhang J, Ma L, Shen W, Shen W, Ma L, Zhang J, Zhang J. The Life Cycle of the Ovary. InOvarian Aging 2023 Aug 15 (pp. 7-33). Singapore: Springer Nature Singapore.
- 17. Parfitt AM. Bone remodelling and bone loss: understanding the pathophysiology of osteoporosis. Clinical obstetrics and gynecology. 1987 Dec 1;30(4):789-811.
- 18. Heaney RP. Pathophysiology of osteoporosis. Endocrinology and metabolism clinics of North America. 1998 Jun 1;27(2):255-65.
- 19. Tayyem R, Abuhijleh H, Al-Khammash A. Lifestyle and Dietary Patterns as Risk Factors for Osteoporosis: A Literature Review. Current Nutrition & Food Science. 2023 Oct 1;19(8):806-16.
- 20. Pisani P, Renna MD, Conversano F, Casciaro E, Di Paola M, Quarta E, Muratore M, Casciaro S. Major osteoporotic fragility fractures: Risk factor updates and societal impact. World journal of orthopedics. 2016 Mar 3;7(3):171.
- 21. McClung MR. The relationship between bone mineral density and fracture risk. Current osteoporosis reports. 2005 Jun;3(2):57-63.
- 22. Holick MF, Dawson-Hughes B, editors. Nutrition and bone health. Totowa NJ:: Humana Press; 2004 May 3.
- 23. El Miedany Y. Bone Healing and Osteoporosis. New Horizons in Osteoporosis Management. 2022:525-45.
- 24. Gold DT. The clinical impact of vertebral fractures: quality of life in women with osteoporosis. Bone. 1996 Mar 1;18(3):S185-9.
- 25. Adams J, Bayes J, Hosseini M, Sibbritt D. Annual out-of-pocket expenditure associated with healthcare use among Australian older women with osteoporosis: a cross-sectional study. Archives of osteoporosis. 2021 Dec;16:1-9.
- 26. Lin LP, Lai WJ, Hsu SW, Lin JD. Health Literacy of Osteoporosis Risks among Caregivers Serving in Disability Care Facilities. International Journal of Environmental Research and Public Health. 2020 Jul;17(13):4903.
- 27. Dennehy C, Tsourounis C. A review of select vitamins and minerals used by postmenopausal women. Maturitas. 2010 Aug 1;66(4):370-80.

- 28. Okiki PA, Osibote IA, Balogun O, Oyinloye BE, Idris OO, Adelegan O, Asoso SO, Olagbemide PT. Evaluation of proximate, minerals, vitamins and phytochemical composition of Moringa oleifera Lam. cultivated in Ado Ekiti, Nigeria. Advances in Biological Research. 2015;9(6):436-43.
- 29. Maizuwo AI, Hassan AS, Momoh H, Muhammad JA. Phytochemical constituents, biological activities, therapeutic potentials and nutritional values of Moringa oleifera (Zogale): a review. Journal of drug design and medicinal chemistry. 2017 Oct 28;3(4):60-6.
- 30. Peñalver R, Martínez-Zamora L, Lorenzo JM, Ros G, Nieto G. Nutritional and antioxidant properties of Moringa oleifera leaves in functional foods. Foods. 2022 Apr 12;11(8):1107.
- 31. Gupta S, Mishra K. Amazing health benefits of Moringa Oliefera and Rosa Canina.
- 32. Fard MT, Arulselvan P, Karthivashan G, Adam SK, Fakurazi S. Bioactive extract from Moringa oleifera inhibits the pro-inflammatory mediators in lipopolysaccharide stimulated macrophages. Pharmacognosy magazine. 2015 Oct;11(Suppl 4):S556.
- 33. Patel C, Rangrez A, Parikh P. The anti-osteoporotic effect of Moringa oliefera on osteoblastic cells: SaOS 2. IOSR J Pharm Biol Sci. 2013 Jan;5(2):10-7.
- 34. FATHIL NM. THE HISTOLOGICAL AND PHYSIOLOGICAL STUDY OF MORINGA OLEIFERA.
- 35. Evelyn SS, Chitra V. Medicinal plants for the treatment of postmenopausal osteoporosis. Biomedical & Pharmacology Journal. 2019;12(3):1561.
- 36. Sun X, Xia T, Zhang S, Zhang J, Xu L, Han T, Xin H. Hops extract and xanthohumol ameliorate bone loss induced by iron overload via activating Akt/GSK3β/Nrf2 pathway. Journal of Bone and Mineral Metabolism. 2022 May;40(3):375-88.
- 37. Ramesh P, Jagadeesan R, Sekaran S, Dhanasekaran A, Vimalraj S. Flavonoids: classification, function, and molecular mechanisms involved in bone remodelling. Frontiers in Endocrinology. 2021 Nov 23;12:779638.
- 38. Stohs SJ, Kaats GR, Preuss HG. Safety and Efficacy of Banaba–Moringa oleifera–Green Coffee Bean Extracts and Vitamin D3 in a Sustained Release Weight Management Supplement. Phytotherapy Research. 2016 Apr;30(4):681-8.
- 39. Garg MK, Kharb S. Dual energy X-ray absorptiometry: Pitfalls in measurement and interpretation of bone mineral density. Indian journal of endocrinology and metabolism. 2013 Mar 1;17(2):203-10.
- 40. Stohs SJ, Kaats GR, Preuss HG. Safety and Efficacy of Banaba–Moringa oleifera–Green Coffee Bean Extracts and Vitamin D3 in a Sustained Release Weight Management Supplement. Phytotherapy Research. 2016 Apr;30(4):681-8.
- 41. Tsang HH, Cheung JP, Wong CK, Cheung PW, Lau CS, Chung HY. Psychometric validation of the EuroQoL 5-dimension (EQ-5D) questionnaire in patients with spondyloarthritis. Arthritis Research & Therapy. 2019 Dec;21:1-4.
- 42. Meireles D, Gomes J, Lopes L, Hinzmann M, Machado J. A review of properties, nutritional and pharmaceutical applications of Moringa oleifera: integrative approach on conventional and traditional Asian medicine. Advances in Traditional Medicine. 2020 Dec;20(4):495-515.
- 43. Chen GL, Xu YB, Wu JL, Li N, Guo MQ. Hypoglycemic and hypolipidemic effects of Moringa oleifera leaves and their functional chemical constituents. Food chemistry. 2020 Dec 15:333:127478.
- 44. Cotabarren J, Claver S, Payrol JA, Garcia-Pardo J, Obregón WD. Purification and characterization of a novel thermostable papain inhibitor from moringa oleifera with antimicrobial and anticoagulant properties. Pharmaceutics. 2021 Apr 8;13(4):512.