



IMPACT OF MEDICATIONS AND DIETARY COMPONENTS ON TYPE II DIABETES TREATMENT: INSIGHTS INTO DRUG-NUTRIENT INTERACTIONS

Dr. Zahra Baig¹, Muhammad Usama², Muhammad Hammad Ul Hassan³, Dr. Saba Ahsan⁴,
Jasmeen Kaur Dhillon⁵, Abdulkadir Abdu⁶, Eaman Hamid⁷, Dr Rashida Bibi^{8*}

¹Demonstrator, Department of Biochemistry, Lahore Medical And Dental College, Pakistan,
Email: zbaig04@gmail.com

²Final Year MBBS Student, Al Nafees Medical College, Islamabad, Pakistan,
Email: osama.alwaha@gmail.com

³Final Year MBBS Student, Al Nafees Medical College, Islamabad, Pakistan,
Email: hamad92qadri@gmail.com

⁴Medical Officer, Department of Internal Medicine, Services Hospital Lahore, Pakistan,
Email: sabakhalid15@gmail.com

⁵MBBS, Primary Health Care Physician, Public Health And Family Welfare Department, Sardar Patel Medical College, Bikaner, India, Email: jk328000@gmail.com

⁶Department of Pharmaceutical Sciences, Gurunanak Dev University Amritsar, Punjab 143005, India, Email: Gaduraabdu61@gmail.com

⁷Doctor, Department of Internal Medicine, Omdurman Islamic university, United States,
Email: hamid3198917@gmail.com

^{8*}School of Nursing and Health, Zhengzhou university, Henan, China,
Email : aallee.zeb@gmail.com

***Corresponding Author:** Dr Rashida Bibi, School of Nursing and Health, Zhengzhou university, Henan, China, Email : aallee.zeb@gmail.com

Abstract:

Background: The prevalence of type II diabetes mellitus (T2DM) is increasing worldwide, particularly among older populations. Long-term management is crucial for clinical improvement, and drug-nutrient interactions may affect treatment efficacy.

Objective: This study aimed to investigate drug-nutrient interactions in T2DM management, focusing on the impact of medications and dietary components on treatment effectiveness.

Methods: An integrated literature review guided the development of a primary question and inclusion/exclusion criteria for selecting relevant studies from PubMed and Google Scholar databases. Six hundred ninety-nine papers were reviewed, summarizing oral medications for DM2 and their modes of action. Additionally, the role of dietary therapy and its susceptibility to drug-nutrient interactions, leading to vitamin deficiencies (e.g., B12) and increased risk of bone fractures, was explored.

Results: Drug-nutrient interactions influence blood sugar regulation and impact patients' nutritional status, including dyslipidemia and weight gain. Certain patient groups, such as women, the elderly, and those with poorly controlled diabetes, are more susceptible to these interactions.

Conclusion: Understanding and routinely assessing drug-nutrient interactions and nutritional status are essential for optimizing T2DM treatment outcomes and enhancing patients' quality of life.

Keywords: Diabetes Mellitus; Nutrients; Interaction; Drugs; Health.

INTRODUCTION:

Hyperglycemia, or elevated blood sugar, is a hallmark of diabetes mellitus (DM), a metabolic condition brought on by malfunctions in the production of or reaction to insulin. As a result, both its incidence and frequency have significantly increased worldwide, particularly in older people. As a result, these figures are rising as a result of both population aging and the longer life expectancy of DM patients (Ahmed et al., 2024; W. Wang et al., 2024).

Since DM causes clinical changes that typically call for ongoing, long-term care, ensuring therapeutic efficacy and reducing potential interference with treatment is critical. Additionally, interactions between pharmaceutical drugs and food or nutritional ingredients in food products may make pharmacological therapies less effective for a variety of illnesses (Wang, Fang, & Yang, 2024; Zahedi et al., 2024).

Additionally, foods designed for specific purposes and providing an appropriate supply of nutrients in separated or mixed form are used exclusively or partly as a supplement to nutrition to aid in the production or control of diabetes mellitus (DM). This approach is based on pharmaceutical and non-pharmacological therapy (Kalva et al., 2024; Zhu et al., 2024).

However, when a drug is consumed with food, modifications may be made to the way it functions or is processed by the body. These modifications may impact the the pharmaceutical or kinetic dynamics of the drug substance and the nutrient (vitamins, minerals), which may affect the therapeutic's efficacy or nutritional balance. This phenomenon is known as a drug-nutrient interaction (Craig et al., 2024; Permana et al., 2024).

Thus, it is imperative to thoroughly examine the recommended diet and the numerous medications prescribed, considering the results of the nutritional assessment and potential interactions with pharmaceutical therapies (Xourafa, Korbmacher, & Roden, 2024).

Thus, this study intends to investigate and evaluate the primary interactions between medications and nutrients during the management of patients with Type II Diabetes Mellitus (DM2), illustrating the primary interactions between drugs and nutrients and how these relationships affect their management and wellness. This is based on a review of the literature (Nurgül & Tülüce, 2024; Peng, 2024).

METHODOLOGY:

This study was carried out in stages as follows and includes an integrated review of the literature: defining the inclusion and exclusion criteria for scientific studies, formulating the main question, searching databases, analyzing study summaries, selecting pertinent studies, assessing the specifics of the selected studies, and analyzing the data gathered (Chawla, Pradhan, & Gupta, 2024; Perakakis et al., 2024; Soto-Mota et al., 2024).

"What are the main interactions among nutrients and drugs in patients with type II diabetes mellitus, and how do these interactions affect the treatment of these patients?" is the study's topic. The "PICO" technique served as the foundation for formulating the guiding research question. In this strategy, "P" stands for patients with type II diabetes mellitus as the population, "I" for the intervention (drug and

nutrition interactions), and "CO" for the setting (interaction evaluation) (Falkenhain et al., 2024; Gregg et al., 2024; Tobias et al., 2024).

To do this, searches in the PubMed and Google Scholar databases were conducted using combinations based on the English terms "Diabetes Mellitus" and "Drug-Nutrient interaction" (Interaction). The study examined free articles published between 2018 and 2023; off-topic, duplicate, paid, and after-hours items were not included (Dimer et al., 2024; He et al., 2024; Ma, 2024).

RESULTS:

The flow diagram in Figure 1 shows that the search yielded 699 articles in the two sources employed (PubMed = 91, Google Scholar = 608) (Pathan et al., 2024; Vajdi et al., 2024).




Figure 1. Flowchart for identification, eligibility criteria, and inclusion of studies in the review.

Table 1 illustrates the scientific papers in the search results according to the study's citation, type, title, and objective (Vajdi et al., 2024; Zhou & Xu, 2024).

QUOTE	TYPE OF STUDY	TITLE	OBJECTIVE
Pereira, 2018	Descriptive investigative research using a numerical methodology.	Drug-nutrient correlations of antihypertensive and antidiabetic medicines prescribed at the Alcides Carneiro University Hospital.	To assess the interactions between antihypertensive and antidiabetic medications and nutrients provided in a university hospital.
Mohn et al., 2018	Literature review	Indications of drug-nutrient interactions when using regularly prescribed drugs for a long time: a report.	Give current information on potential drug-nutrient interactions, concentrating on the most often given drugs for diseases usually diagnosed in the adult US population.

Owen et al., 2021	Literature review	Metformin, folate, and vitamin B12 interactions and their possible effects on fetal development and long-term metabolic health in diabetic pregnancies.	Examine how metformin, folate, and vitamin B12 interact throughout a diabetic woman's pregnancy and how this may impact the fetus's development and long-term metabolic health.
Dantas, 2019	Literature review	Drug-food interaction in older adults with type 2 diabetes mellitus.	To examine the relationship between food and medication in older adults with type 2 diabetes mellitus to find potential barriers to treatment efficacy.
Leal and Júnior, 2018	Literature review	Drug-nutrient interaction: description and novel approaches to assessment.	Examine how dietary supplements and medications interact and how this may impact the efficacy of medical interventions.

Table 1. Selected articles, citations, type of study, title, and objective.

Table 2 shows the results of a survey conducted in addition to the data shown in Table 1 and includes the primary classes of oral medications used to treat DM2. The drug label and mechanism of action were then listed (Frías et al., 2024; TAHA, Abd El-Azeaz, & ABD EL-RAZIK, 2025).

CLASS	MEDICINES	ACTION
BIGUANIDES	Metformin	It lessens the quantity of glucose that the liver releases. Increases susceptibility to insulin's effects.
	Metformin XR	
SULFONYLUREAS	Chlorpropamide	Causes the pancreas to produce more insulin when stimulated.
	Glibenclamide	
	Glipizide	
	Gliclazide	
	Gliclazide MR	
	Glimepiride	
MEGLITINIDES	Repaglinide	It causes an increase in insulin secretion by the pancreas right after a meal.
	Nateglinide	
GLITAZONES	Pioglitazone	Increases the production and release of insulin.
	Thiazolidinediones	
ALPHA-GLUCOSIDASE INHIBITORS	Acarbose	Slows down the process of absorbing carbs.
GLIPTINS	Sitagliptin	Increases the production and release of insulin.
	Vildagliptin	
	Linagliptin	
	Alogliptin	
SGLT2 INHIBITORS	Dapagliflozin	It increases the amount of extra glucose excreted in the urine by preventing the kidneys from reabsorbing glucose.
	Empagliflozin	

Table 2. Main classes of drugs used in the treatment of type II diabetes mellitus.

DISCUSSION:

As defined by the Institute of Medicine (IMS), dietary therapy treats a disease or condition by altering the amount of nutrients consumed. As a result, it is critical to attain appropriate glucose control, a crucial element in the effectiveness of DM2 medication therapy (Mohammadi et al., 2024; Paul et al., 2024).

As a result, the relationship between nutrition and medications has received increasing attention, maybe due to medications becoming more precise and effective. Furthermore, drugs and foods have numerous common traits, including physical aspects that can influence some biological responses and other dose-related adverse consequences (Hussain, Borah, & Ahmed, 2024; Tolentino & Brynes, 2024).

Pereira's research highlighted that metformin, a medication commonly prescribed as the first treatment for diabetes mellitus, needs particular attention since it can interfere with the metabolism of vitamin B12. This study and the one by Mohn and associates about metformin use can be correlated. Metformin use may lower vitamin B12 levels because of potential issues with intestinal absorption (Ghahfarrokhi et al., 2024; Wood & Pratt, 2024).

Thus, when taking the medication, those who have diabetes or are more likely to have low vitamin B12 levels may be particularly sensitive to these side effects. It was also noted in the research by Mohn and associates that extended usage of thiazolidinediones raises the likelihood of bone fractures. Furthermore, it has been shown that people with T2DM receiving antidiabetic medicine do not consume enough calcium, vitamin D, or magnesium nutrients necessary for maintaining healthy bones (Bonekamp et al., 2024; Manzano, Llames, & Macabeo, 2024).

According to Owen and colleagues' study, the action of metformin is similar to that of antifolates and can lower vitamin B12 levels. This can have long-term adverse effects, particularly for pregnant women with DM2 and gestational diabetes mellitus. (DMG), compromising aerobic respiration's mitochondrial activity and carbon metabolism (CG, Adibe, Ukwe, & Aguwa, 2030).

Dantas claims his research looked at potential interactions between medications and meals high in folic acid and vitamin B12. With a rate of 35.5%, metformin was used more frequently than glimepiride, alogliptin and metformin, dapagliflozin and metformin, glimepiride, alogliptin and metformin, and metformin, which was used less frequently with a rate of just 1.5%. Since of this interaction, the body absorbs fewer vitamins, which can be detrimental to older persons since it increases the likelihood that they will become deficient in these vitamins and develop other health issues linked to nutrient deficiencies (Gierlach et al., 2024; Khairnar et al., 2024).

Moreover, the quality of life for senior DM2 patients and the efficacy of medication therapy may be affected by this interaction. In this context, metformin use remained to demonstrate a substantial connection with B12 deficiency even after accounting for characteristics like age, body mass index, use of vitamin B12 supplements, and medications that lower gastric acidity (Patil & Tupe, 2024; Thompson et al., 2024).

According to Leal and Júnior's study, the effects of drugs on nutritional status can be complex because they can affect food intake, digestion, absorption, and nutritional metabolic functioning. These changes in metabolic function can result from drug use and include weight gain, hyperglycemia, dyslipidemia, and other conditions. Furthermore, a correlation between weight gain and elevated blood glucose levels linked to medication use has been noted in certain studies, which may present further challenges for the management of diabetes mellitus (Salh, Aziz, Ahmed, & Mahwi, 2024).

As a result, specific patient populations such as women, the elderly, those who consume too many calories, and those with poorly managed diabetes are more likely to experience medication interactions (Shao, Shi, Nauman, Price-Haywood, & Stoecker, 2024).

CONCLUSION:

Consequently, we can conclude that drug-nutrient interactions are essential when treating DM2 patients based on the data in the study. Moreover, these interactions may affect patients' nutritional state and the efficacy of their medication regimen.

Moreover, it has been noted that metformin, a medication frequently prescribed for the first course of therapy for diabetes mellitus, may have adverse effects on the body's ability to absorb vitamin B12,

potentially leaving patients deficient in it. In this regard, people with DM2 have been linked to an elevated incidence of bone fractures due to various medications, including thiazolidinediones. It has also been discovered that patients with DM2 receiving antidiabetic medication do not consume enough calcium, vitamin D, or magnesium, which might harm bone health. It is crucial to remember that not every patient is affected by these interactions similarly and may differ depending on age, body mass index, and supplement use. To guarantee efficient treatment and the preservation of a patient's quality of life, it is crucial to be aware of these correlations and to evaluate the nutritional state of the patients periodically.

REFERENCES:

1. Ahmed, S., Ripon, M. S. H., Islam, M. F., Ullah, A., Sultan, S., Sajid, M., & Rahman, T. (2024). Association of dietary intake and nutrition knowledge with diabetes self-management behavior among Bangladeshi type 2 diabetes mellitus adults: A multi-center cross-sectional study. *Endocrine and Metabolic Science*, 14, 100156.
2. Bonekamp, N. E., Visseren, F. L., Cramer, M. J., Dorresteijn, J. A., van der Meer, M. G., Ruigrok, Y. M., . . . Koopal, C. (2024). Long-term lifestyle change and risk of mortality and Type 2 diabetes in patients with cardiovascular disease. *European Journal of Preventive Cardiology*, 31(2), 205-213.
3. CG, A.-O., Adibe, M., Ukwe, C., & Aguwa, C. (2030). *Journal of Pharmaceutics and Therapeutics* The Impact of Family Support on Medication Adherence and Glycemic Control of Type 2 Diabetes Outpatients in a Nigerian Tertiary Hospital.
4. Chawla, G., Pradhan, T., & Gupta, O. (2024). An insight into the combat strategies for treating type 2 diabetes mellitus. *Mini-Reviews in Medicinal Chemistry*, 24(4), 403-430.
5. Craig, H., Le Roux, C., Alsaeed, D., Al-Ozairi, E., Heneghan, H., & Al-Najim, W. (2024). Factors that determine patients considering medication for the disease of obesity: an IMI2 SOPHIA study.
6. Dimer, L. M., Colonetti, T., Uggioni, M. L. R., Ferraz, S. D., Grande, A. J., da Rosa, M. I., & Ceretta, L. B. (2024). Evaluating the effects of *Bauhinia forficata* as an adjuvant in treating type 2 diabetes mellitus: systematic review and meta-analysis. *European Journal of Integrative Medicine*, 102331.
7. Falkenhain, K., Oliveira, B. F., Islam, H., Neudorf, H., Cen, H. H., Johnson, J. D., . . . Little, J. P. (2024). The effect of acute and 14-day exogenous ketone supplementation on glycemic control in adults with type 2 diabetes: two randomized controlled trials. *American Journal of Physiology-Endocrinology and Metabolism*, 326(1), E61-E72.
8. Frías, J. P., Galindo, R. J., Wang, H., Malik, R. E., Chivukula, K. K., & Maldonado, J. M. (2024). Tirzepatide in Hispanic/Latino patients with type 2 diabetes: a subgroup analysis of the SURPASS Program. *The Journal of Clinical Endocrinology & Metabolism*, 109(2), 557-568.
9. Ghahfarrokhi, M. M., Shirvani, H., Rahimi, M., Bazgir, B., Shamsadini, A., & Sobhani, V. (2024). Feasibility and preliminary efficacy of different intensities of functional training in elderly type 2 diabetes patients with cognitive impairment: a pilot randomized controlled trial. *BMC geriatrics*, 24(1), 1-15.
10. Gierlach, K., Nowak, K., Olejarz, Z., Słowik, M., Drygała, Z., Wyrwał, J., . . . Krasuska, M. (2024). Critical Analysis of the Efficacy and Limitations of GLP-1 Pharmaceuticals in Obesity Management: A Comprehensive Literature Review. *Journal of Education, Health, and Sport*, 55, 87-103.
11. Gregg, E. W., Chen, H., Bancks, M. P., Manalac, R., Maruthur, N., Munshi, M., . . . Group, L. A. R. (2024). Impact of remission from type 2 diabetes on long-term health outcomes: the Look AHEAD study findings. *Diabetologia*, 1-11.
12. He, A., Shi, C., Wu, X., Sheng, Y., Zhu, X., Yang, J., & Zhou, Y. (2024). Clusters of Body Fat and Nutritional Parameters are Strongly Associated with Diabetic Kidney Disease in Adults with Type 2 Diabetes. *Diabetes Therapy*, 15(1), 201-214.

13. Hussain, Z., Borah, M. D., & Ahmed, R. K. (2024). Computational methods for studying the relationship between nutritional status and respiratory viral diseases: a systematic review. *Artificial Intelligence Review*, 57(1), 3.
14. Kalva, P., Shi, A., Kakkilaya, A., Saleh, I., Albadour, M., & Kooner, K. (2024). Associations between depression and diabetic retinopathy in the National Health and Nutrition Examination Survey, 2011 to 2018. Paper presented at the Baylor University Medical Center Proceedings.
15. Khairnar, S. J., Ahire, E. D., Jagtap, M. R., Surana, K. R., Kshirsagar, S. J., & Keservani, R. K. (2024). Management and Prevention of Diseases by Flavonoids Advances in Flavonoids for Human Health and Prevention of Diseases (pp. 47-71): Apple Academic Press.
16. Ma, S. (2024). Dietary nutrition treatment for older people with hyperglycemia. Paper presented at the Third International Conference on Biological Engineering and Medical Science (ICBioMed2023).
17. Manzano, J. A. H., Llamas, L. C. J., & Macabeo, A. P. G. (2024). Tetrahydrobisbenzylisoquinoline alkaloids from *Phaeanthus ophthalmics* inhibit target enzymes associated with type 2 diabetes and obesity. *Journal of Applied Pharmaceutical Science*, 14(1), 230-237.
18. Mohammadi, F., Yadegar, A., Rabizadeh, S., Ayati, A., Seyedi, S. A., Nabipoorashrafi, S. A., . . . Nakhjavani, M. (2024). Correlates of ordinary and decreased HDL cholesterol levels in type 2 diabetes: a cohort-based cross-sectional study. *Lipids in Health and Disease*, 23(1), 1-13.
19. Nurgül, G., & Tülüce, D. (2024). Diabetes Attitude and Self-Efficiency Levels of Patients with Type 2 Diabetes Mellitus Taking Insulin. *Bilecik Şeyh Edebali Üniversitesi Sağlık Bilimleri Fakültesi Dergisi*, 2(1), 37-52.
20. Pathan, A. S., Ahire, M. R., Diwane, S. A., Jain, P. G., Pandagale, P. M., & Ahire, E. D. (2024). Functional Foods in Prevention of Diabetes Mellitus Applications of Functional Foods in Disease Prevention (pp. 139-164): Apple Academic Press.
21. Patil, R. S., & Tupe, R. S. (2024). Communal interaction of glycation and gut microbes in diabetes mellitus, Alzheimer's, and Parkinson's disease pathogenesis. *Medicinal Research Reviews*, 44(1), 365-405.
22. Paul, T., Pathak, K., Saikia, R., Gogoi, U., Jyoti Sahariah, J., & Das, A. (2024). The Role of Medicinal Plants in the Treatment and Management of Type 2 Diabetes. *Current Traditional Medicine*, 10(2), 83-96.
23. Peng, Y. (2024). A meta-analysis of drugs benefiting diabetic patients by modulating gut flora. Paper presented at the Third International Conference on Biological Engineering and Medical Science (ICBioMed2023).
24. Perakakis, N., Bornstein, S. R., Birkenfeld, A. L., Linkermann, A., Demir, M., Anker, S. D., . . . Ruilope, L. M. (2024). Efficacy of finerenone in patients with type 2 diabetes, chronic kidney disease and altered markers of liver steatosis and fibrosis: A FIDELITY subgroup analysis. *Diabetes, Obesity and Metabolism*, 26(1), 191-200.
25. Permana, H., Soetedjo, N. N. M., Yanto, T. A., Tendean, M., Hariyanto, T. I., & Suastika, K. (2024). Different doses of imeglimin for management of type 2 diabetes mellitus: a systematic review, meta-analysis, and meta-regression of randomized clinical trials. *Expert Review of Endocrinology & Metabolism*, 19(1), 89-98.
26. Salh, H. J. H., Aziz, T. A., Ahmed, Z. A., & Mahwi, T. O. (2024). Association between Albuminuria, Glycated Hemoglobin with Comorbidities in Type 2 Diabetes Patients: Experience in Sulaimani City, Iraq. *Al-Rafidain Journal of Medical Sciences (ISSN 2789-3219)*, 6(1), 1-8.
27. Shao, Y., Shi, L., Nauman, E., Price-Haywood, E., & Stoecker, C. (2024). Telehealth use and its impact on clinical outcomes in patients with type 2 diabetes during the COVID-19 pandemic. *Diabetes, Obesity and Metabolism*, 26(1), 118-125.
28. Soto-Mota, A., Flores-Jurado, Y., Norwitz, N. G., Feldman, D., Pereira, M. A., Danaei, G., & Ludwig, D. S. (2024). Increased LDL-cholesterol on a low-carbohydrate diet in adults with typical but not high body weight: a meta-analysis. *The American Journal of Clinical Nutrition*.

29. TAHA, N. M., Abd El-Azeaz, M., & ABD EL-RAZIK, B. G. (2025). Factors affecting compliance of diabetic patients toward therapeutic management. *Diabetes*, 5, 6.
30. Thompson, A. S., Candussi, C. J., Tresserra-Rimbau, A., Jennings, A., Bondonno, N. P., Hill, C., . . . Kühn, T. (2024). A healthful plant-based diet is associated with lower type 2 diabetes risk via improved metabolic state and organ function: A prospective cohort study. *Diabetes & Metabolism*, 50(1), 101499.
31. Tobias, D. K., Hamaya, R., Clish, C. B., Liang, L., Deik, A., Dennis, C., . . . Manson, J. E. (2024). Type 2 diabetes metabolomics score and risk of progression to type 2 diabetes among women with a history of gestational diabetes mellitus. *Diabetes/Metabolism Research and Reviews*, 40(1), e3763.
32. Tolentino, D. A., & Brynes, M. E. (2024). Filipino Americans' Social and Cultural Experiences of Type 2 Diabetes Management: Cultural Paradox, Ownership, and Success Definition. *Journal of Transcultural Nursing*, 35(1), 41-52.
33. Vajdi, M., Seyedhosseini-Ghaheh, H., Hassanizadeh, S., Khajeh, M., Tabrizi, F. P. F., & Askari, G. (2024). Effect of sesame supplementation on body composition, and lipid profile in patients with type 2 diabetes: A Systematic Review and Meta-analysis of randomized controlled trials. *Nutrition, Metabolism and Cardiovascular Diseases*.
34. Wang, W., Li, X., Chen, F., Wei, R., Chen, Z., Li, J., . . . Guo, L. (2024). Secondary analysis of newly diagnosed type 2 diabetes subgroups and treatment responses in the MARCH cohort. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 18(1), 102936.
35. Wang, X., Fang, J., & Yang, L. (2024). Research progress on ocular complications caused by type 2 diabetes mellitus and the function of tears and blepharons. *Open Life Sciences*, 19(1), 20220773.
36. Wood, A. C., & Pratt, K. (2024). School-based health: an opportunity to address challenges in managing pediatric obesity and type 2 diabetes. *Current Opinion in Pediatrics*, 36(1), 33-41.
37. Xourafa, G., Korbmacher, M., & Roden, M. (2024). Inter-organ crosstalk during development and progression of type 2 diabetes mellitus. *Nature Reviews Endocrinology*, 20(1), 27-49.
38. Zahedi, M., Motahari, M. M., Fakhri, F., Aphshari, N. M., Poursharif, S., Jahed, R., & Nikpayam, O. (2024). Is vitamin D deficiency associated with retinopathy in type 2 diabetes mellitus? A case-control study. *Clinical Nutrition ESPEN*, 59, 158-161.
39. Zhou, X., & Xu, J. (2024). Association between serum uric acid-to-high-density lipoprotein cholesterol ratio and insulin resistance in patients with type 2 diabetes mellitus. *Journal of Diabetes Investigation*, 15(1), 113-120.
40. Zhu, K., Qian, F., Lu, Q., Li, R., Qiu, Z., Li, L., . . . Yang, K. (2024). Modifiable Lifestyle Factors, Genetic Risk, and Incident Peripheral Artery Disease Among Individuals With Type 2 Diabetes: A Prospective Study. *Diabetes Care*, dc231503.