



An investigation to assess the implant removal for a variety of reasons

1 Dr Hasnain Ali

Resident orthopedics, Shifa international hospital, Islamabad

2Dr Zofeen Aftab

2BHU city Mzd

3Afan Shamim

3Bolan Medical Complex Hospital

4Dr. Muhammad Umair Khan

4Dr. Sikandar Ali Mandhro Civil Hospital Badin, Managed by The Indus Hospital and Health Network Karachi, Sindh, Pakistan

5Dr Farnaz Ali

Medical officer DHQ Bhimber.

6Memoona Fareed

6H.H. SKBZH Muzaffarabad

7Kashif Lodhi

7Department of Agricultural, Food and Environmental Sciences. Università Politénica delle Marche Via Brece Bianche 10, 60131 Ancona (AN) Italy,

ABSTRACT:

INTRODUCTION: As technology has improved, fracture treatment has as well. Modern contouring plates and nails are possible depending on the bone and the location of the fracture. even after implant material advances. On the other side, there is an ongoing debate among surgeons over the crucial justification for implant removal. The patient's symptoms are mostly responsible for implant removal. The study's objectives are to examine the main factors that lead to implant removal and identify the implants that must be removed during orthopedic procedures.

METHODS: An analysis of 89 patients who were hospitalized for implant removal between January 2017 and January 2020 was carried out. A total of 112 patients were selected with care. Selected were all implant removal patients who were hospitalized with ages ranging from 18 to 55 and who had all necessary paperwork in order, such as a completed blood test and an X-ray report, at the time of admission.

RESULTS: 89 patients participated in the trial, including 73 men and 16 women. The research group's average age was 38 years old. Four categories were used to group the reasons for implant removal: (I) Pain, (II) Infection, (III) Implant resorption or failure (IV) Opt-in (patient's decision). According to our study, distal Tibial/Ankle plates (14.4%), Femoral Intramedullary (IMIL) nails (13.25%), Olecranon wires and

plates (12.04%), Tibial IMIL nails, and Patellar tension band wires (9.53%) are implants that are commonly removed.

CONCLUSIONS: The medical justifications for implant removal have not been well investigated. Our investigation revealed that pain or discomfort was the main reason for implant removal (37.1%). Infection was the second most frequent cause of removal (26.96%) of the implant. Thirdly, implant failure in symptomatic patients due to subpar implants, insufficient or defective implants, and unwilling patients was an indicator. The patient's desire was the fourth indication. Due to variables including bone ingrowth and implant deterioration, implant removal requires a difficult surgical procedure.

KEYWORDS: fracture, implant, nails, plates

INTRODUCTION: With the advancement of technology, fracture treatment has also become better. To support the bone and encourage correct healing, fracture care with implants entails the use of surgical hardware, such as plates, screws, nails, rods, and pins. These implants may be removed after the bone has completely healed and are used to keep the bone in the right position as it heals. Long bones of the body like the femur, tibia, and humerus are often treated for fractures using plates and screws. The thighbones (femur) and shinbones (tibia) are often treated with intramedullary nails, but the hand, foot, and ankle bones may be treated with external fixators, which are devices put to the outside of the skin and linked to the bone using pins or wires.

Implants come in a variety of forms, including bioabsorbable, metallic, and hybrid. The surgeon will choose the best fracture treatment option based on the kind of fracture. Modern contour plates and nails are available depending on the bone and the location of the fracture. In this century, there are several alternatives for surgical fracture care due to development. [1,2] Particularly in the design of the implants, such as Intramedullary (IM) nails, plates, and screws specifically for unusual bone frames and boosting rate of union and treatment effectiveness. Stainless steel and titanium alloys are the most often utilized materials in the production of orthopedic implants. [3,4] Due to its lower weight and infection resistance. [5] In rare circumstances, implants that have served their function may still be recognized by the body as a foreign object and cause a response.

It is important to remember that complications with implants, such as infection, non-union, or improper union, are possible. So it's essential to provide the right treatment and follow-up. After surgery, the patient often needs physical therapy and rehabilitation to help them restore their strength and range of motion.

Due to the lack of evidence-based recommendations, there is still significant debate among surgeons worldwide about the removal of asymptomatic implants. [6,7] Children's implants are often removed when their fractured bones have fused, as per usual procedure, to avoid interfering with growth plates or causing additional infections, corrosion, or reactions to foreign bodies. [8] The two main reasons to remove implants are to relieve discomfort and enhance functionality. The technique is a taxing operation that carries a risk of refractures or neurovascular damage. [9-12] The key reasons for implant removal will be surveyed in this research, along with the implants that need to be removed during orthopedic procedures.

METHODS: 89 individuals who were hospitalized for implant removal from December 2019 to December 2022 were the subject of this study. A total of 112 patients were selected with care. A list of inclusion criteria was used to choose the patients. Adult patients between the ages of 18 and 55 who were hospitalized for implant removal in the orthopedic department's wards had to meet the inclusion criteria. They also had to have completed all necessary investigations, such as basic blood work and an x-ray of the implant removal location, at admission the time. Written records of the surgery's obtained consent, post-operative care, and discharge summaries are required. The outcomes and follow-ups after the study group was eliminated should also be recorded. Exclusion criteria for children under the age of 16 included past surgeries including K-wire fixation, external fixators in situ, cases involving joint prostheses in situ, and loss of substantial documented data. Due to inadequate data, other conditions including diabetes, and inconsistent medication use, 23 instances were ruled out.

RESULTS: Seventy-three (82%) of the 89 patients were males, and 16 (8%) were women. Although they ranged in age from 18 to 55, their average age was 38. Four kinds of primary reasons for implant removal

were identified: (I) Pain or discomfort, (II) infection, (III) implant failure or resorption, and (IV) elective surgery of the patient's choosing. [Table 1].

Table 1: Cases distribution

Implant type	Prominent	Infected	Implant	Elective	Other reasons
Ankle implants/ distal tibial	2	8	2		
Tibial plates	1	1			
Tibial nails		2		6	
Proximal tibial plates		4		2	
Patella TBW	8				
Femoral plates	2			2	
Femoral nails	4	2	3	2	
Hip plates and screws	1	1	2		
Forearm plates				3	1
Olecranon TBW/plates	7	3			
Distal humeral plates	6			2	
Humeral diaphysis plate/nail	2	2	1		

Table 2: AO recommendations regarding when to remove implants after simple fracture healing


Bone Fracture	Time after implantation (Months)
Upper extremity	12 to 18
Pelvis	From 10th day of month
Peritrochanteric and femoral neck fractures	12 to 18
Femoral intramedullary (IM) nail	24 to 36
Femoral shaft double plates	18 to 24
Femoral shaft single plate	24 to 36
Femoral condyles	12 to 24
Patella TBW	8 to 12
Proximal tibia	12 to 18
Tibial intramedullary (IM) nail	18 to 24
Tibial shaft plate	12 to 18
Tibial pilon	12 to 18
Malleolar fractures	8 to 12

(I). Pain, discomfort, or predominance was seen in 33 patients (37.1%). The average time after fracture fixation was 38 months, however, it ranged from 6 to 84 months. According to our study, the most often removed implants were the femoral IM nail, the patella tension band wire (TBW), the olecranon TBW/plates, and the twelve. The average length of stay for the patients in the study was 7 days, and after six consecutive months of follow-ups, the results indicated that 30 out of 33 patients had eliminated their pain (90.9%), while the other 3 patients continued to have slight pain or discomfort (9.09%).

(II) 24 patients (26.96%) needed implant removal due to infected implants already in situ. The period after fracture fixation varied from 2 to 56 months, or 47.57 months on average. While the contaminated implants were being removed from the other 23 patients, the fracture of one patient, which didn't seem to be connected, was treated further using external fixators. The most often removed implants, according to our

An investigation to assess the implant removal for a variety of reasons

data, were olecranon plates (4%), proximal tibial plates (9%) and distal tibial/ankle plates and screws (11). The average length of stay for the patients in the study was 7 days, and the results indicated that 21 out of 24 patients (87.5%) had fully recovered from infection by the time they were discharged, whereas 3 patients had developed chronic osteomyelitis. [Figure 1 (a, b), Chart 2].



Chart



Figure 1: Implant removal

(III). Due to implant failure, 8 patients (8.98%) had revision osteosynthesis in addition to implant removal. The average time since the first operation was 2 to 12 months. Our study found that humeral shaft dynamic compression plates (1), femoral IMIL nails (2), distal tibial plates (3), cannulated cancellous screws in the femoral neck (2), and cannulated cancellous screws in the tibia (1) were the implants that were removed the most often. Following fixing surgery, a follow-up exam indicated significant bone loss beneath both forearm plating bones [Figure 1 (d)]. Both plates were eventually removed. There were no new problems with additional follow-up.

(IV). Despite having no symptoms, 24 patients (26.97%) decided to have their implants removed on their own [Chart 4].

According to a further study of the data, no implants were removed following any serious vascular damage or refracture. After having the distal humeral plates removed, a patient had ulnar nerve palsy; nevertheless, they improved with further physiotherapy treatment. Chronic osteomyelitis caused by an infected Tibial IMIL nail was treated with sequestrectomy and wound washing.

DISCUSSIONS: The purpose of the research was to identify the primary reasons for implant removal from orthopedic operations as well as which implants needed to be removed. The study included the records of 89 adult patients who were hospitalised for implant removal and the records of 112 adult patients from Mayo Hospital in Lahore.

The AO association advises prompt implant removal. [13,14] The right clinical justifications for implant removal have not been established. As a result of variables including bone engraftment and wear on the implants, implant evacuation is a difficult procedure that carries a risk of refractures or neurovascular damage. [15,16,17]

The research revealed a male predominance, while 82% of implant removal patients were male. Similar results were reported in second research, which showed a male predominance (189 out of 275 patients), and in a third study, which showed that 30 out of 40 patients (or 75%) were male. [18]

The bulk of implant removal operations seem to be carried out on male patients. The study would have been far more accurate if there had been an equal number of male and female patients. In our research, 30 of 33 patients had total pain alleviation (90.9%), while the last three patients continued to have slight pain or discomfort (9.09%), which was also seen in previous studies. [19,20] In a different research, 10 (20%) of the 51 asymptomatic individuals who had implants removed had problems. [21-25] By our research, pain or discomfort was the primary factor (37.1%) in implant removal. Other studies have found comparable results. [26,27,28] Infection was the second most frequent sign that the implant had to be removed (26.96%). [29] According to research, infections may develop in 5% of all implant removals. [30] Thirdly, symptomatic patients with implant failure due to poor implants, inadequate or flawed implants, or uncooperative patients was an indication. Three more research had similar findings. [31,32] The patient's desire was the fourth indication.

CONCLUSIONS: In certain circumstances, implant removal may be required, such as when the patient is dealing with persistent discomfort, implant protrusion, or infection. Removing an implant may also be necessary due to patient demand, implant failure, or bone resorption. It may be especially difficult to remove weight-bearing implants without causing more injury since they may have fused to the surrounding bone. The total elimination of symptoms cannot be ensured by removing the implant. The bone and surrounding tissue may sustain more damage as a result of the implant removal, necessitating additional surgery or rehabilitation. Even after the implant has been removed, some individuals may still feel pain or other symptoms. Therefore, before opting to remove the implant, it should be carefully thought out and evaluated together with the patient's health and available treatments.

REFERENCES:

1. Chung, P. H., Leong, J. Y., Phillips, C. D., & Henry, G. D. (2022). Microorganism profiles of penile prosthesis removed for infection, erosion, and mechanical malfunction based on next-generation sequencing. *The Journal of Sexual Medicine*, 19(2), 356-363.
2. Kreutzer, K., Steffen, C., Nahles, S., Koerdt, S., Heiland, M., Rendenbach, C., & Beck-Broichsitter, B. (2022). Removal of patient-specific reconstruction plates after mandible reconstruction with a

- fibula free flap: is the plate the problem?. *International journal of oral and maxillofacial surgery*, 51(2), 182-190.
3. Tas, D. B., Smeeing, D. P., Keizer, J., Houwert, R. M., & Emmink, B. L. (2022). Postoperative complications of minimally invasive intramedullary nail fixation versus plate fixation for distal fibular fractures in elderly patients: a retrospective double cohort study in a geriatric trauma unit in the Netherlands. *The Journal of Foot and Ankle Surgery*, 61(6), 1170-1176.
 4. Kweh, B. T. S., Tan, T., Lee, H. Q., Hunn, M., Liew, S., & Tee, J. W. (2022). Implant removal versus implant retention following posterior surgical stabilization of thoracolumbar burst fractures: A systematic review and meta-analysis. *Global Spine Journal*, 12(4), 700-718.
 5. Tay, M. L., McGlashan, S. R., Monk, A. P., & Young, S. W. (2022). Revision indications for medial unicompartamental knee arthroplasty: a systematic review. *Archives of Orthopaedic and Trauma Surgery*, 1-14.
 6. Bicong, L., Allen, J. C., Arps, K., Al-Khatib, S. M., Bahnson, T. D., Daubert, J. P., ... & Piccini, J. P. (2022). Leadless pacemaker implantation after lead extraction for cardiac implanted electronic device infection. *Journal of cardiovascular electrophysiology*, 33(3), 464-470.
 7. Casiraghi, A., Galante, C., Rohayem, M., Vittone, G., Domenicucci, M., Cattaneo, S., ... & Milano, G. (2023). Implant retention with serial debridement and use of antibiotic-loaded calcium sulphate beads in acute fracture-related infection (FRI) after pelvic ring or acetabular fractures: a retrospective case series of 7 cases. *Injury*.
 8. Rizaev, J. A., & Bekmuratov, L. R. (2022). Prevention of tissue resorption during immediate implant placement by using socket shield technique. *Art of Medicine. International Medical Scientific Journal*, 2(3).
 9. Lin, Y. C., Chang, C. H., Hu, C. C., Chang, Y., Lee, S. H., & Lin, S. H. (2023). Does the indication for revision associate with the outcomes of revision total knee arthroplasty? A retrospective analysis with a minimum ten-year follow-up.
 10. Wixtrom, R., Glicksman, C., Kadin, M., Lawrence, M., Haws, M., Ferenz, S., ... & McGuire, P. (2022). Heavy metals in breast implant capsules and breast tissue: findings from the systemic symptoms in women–biospecimen analysis study: part 2. *Aesthetic Surgery Journal*, 42(9), 1067-1076.
 11. Tułecki, Ł., Czajkowski, M., Targońska, S., Polewczyk, A., Jacheć, W., Tomków, K., ... & Kutarski, A. (2022). The role of cardiac surgeon in transvenous lead extraction: Experience from 3462 procedures. *Journal of Cardiovascular Electrophysiology*, 33(7), 1357-1365.
 12. Mavrommatis, S., LaRoque, M. C., Yang, G., Brahme, I. S., & Cole, P. A. (2022). Does the Option to Keep Explanted Orthopaedic Trauma Implants Influence Patient Satisfaction and Perception of Care? Results of a Survey Study. *Journal of Orthopaedic Trauma*, 10-1097.
 13. Diemberger, I., Segreti, L., Rinaldi, C. A., Svendsen, J. H., Kutarski, A., Younis, A., ... & ELECTRa Investigators. (2022). Transvenous Lead Extraction in Patients with Cardiac Implantable Device: The Impact of Systemic and Local Infection on Clinical Outcomes—An ESC-EHRA ELECTRa (European Lead Extraction Controlled) Registry Substudy. *Biology*, 11(4), 615.
 14. Diemberger, I., Segreti, L., Rinaldi, C. A., Svendsen, J. H., Kutarski, A., Younis, A., ... & ELECTRa Investigators. (2022). Transvenous Lead Extraction in Patients with Cardiac Implantable Device: The Impact of Systemic and Local Infection on Clinical Outcomes—An ESC-EHRA ELECTRa (European Lead Extraction Controlled) Registry Substudy. *Biology*, 11(4), 615.
 15. Ravidà, A., Siqueira, R., Di Gianfilippo, R., Kaur, G., Giannobile, A., Galindo-Moreno, P., ... & Wang, H. L. (2022). Prognostic factors associated with implant loss, disease progression or favorable outcomes after peri-implantitis surgical therapy. *Clinical Implant Dentistry and Related Research*, 24(2), 222-232.
 16. Wilson, S. K., & Picazo, A. L. L. (2022). Update on the Penuma® an FDA-cleared penile implant for aesthetic enhancement of the flaccid penis. *International Journal of Impotence Research*, 34(4), 369-374.

17. Sorrentino, F., Tealdo, G., Cazzador, D., Favaretto, N., Brotto, D., Montino, S., ... & Zanoletti, E. (2022). Cochlear implant in vestibular schwannomas: long-term outcomes and critical analysis of indications. *European Archives of Oto-Rhino-Laryngology*, 1-10.
18. Patel, S., Ghosh, A., Jindal, K., Kumar, V., Aggarwal, S., & Kumar, P. (2022). Spinopelvic fixation for vertically unstable AO type C pelvic fractures and sacral fractures with spinopelvic dissociation-A systematic review and pooled analysis involving 479 patients. *Journal of Orthopaedics*.
19. Schwarz, F., Obreja, K., Mayer, S., Ramanauskaitė, A., Sader, R., & Parvini, P. (2022). Efficacy of autogenous tooth roots for a combined vertical and horizontal alveolar ridge augmentation and staged implant placement. A prospective controlled clinical study. *Journal of Clinical Periodontology*, 49(5), 496-505.
20. Chang, C. J., Huang, T. C., Hoshino, Y., Wang, C. H., Kuan, F. C., Su, W. R., & Hong, C. K. (2022). Functional outcomes and subsequent surgical procedures after arthroscopic suture versus screw fixation for ACL tibial avulsion fractures: a systematic review and meta-analysis. *Orthopaedic Journal of Sports Medicine*, 10(4), 23259671221085945.
21. Sun, Y., Helmholz, H., Will, O., Damm, T., Wiese, B., Luczak, M., ... & Willumeit-Römer, R. (2022). Dynamic in vivo monitoring of fracture healing process in response to magnesium implant with multimodal imaging: Pilot longitudinal study in a rat external fixation model. *Biomaterials Science*, 10(6), 1532-1543.
22. Stapińska-Syniec, A., Sobstyl, M., & Paskal, W. (2023). Skin-related complications following deep brain stimulation surgery: A single-center retrospective analysis of 525 patients who underwent DBS surgery. *Clinical Neurology and Neurosurgery*, 225, 107571.
23. van Sloten, M., Gómez-Junyent, J., Ferry, T., Rossi, N., Petersdorf, S., Lange, J., ... & ESCMID Study Group of Implant Associated Infections (ESGIAI). (2022). Should all patients with a culture-negative periprosthetic joint infection be treated with antibiotics? A multicentre observational study. *The Bone & Joint Journal*, 104(1), 183-188.
24. de Carvalho, L. F., de Carvalho, L. P., Sotto-Maior, B. S., Dias, A. L., Bezerra, F. J. B., Bergamo, E. T., & de Carvalho, A. M. (2022). Rehabilitation of Atrophic Maxilla With Immediate Loading of Extrasinus Zygomatic Implant. *Journal of Craniofacial Surgery*, 33(5), e488-e491.
25. Badin, D., Ortiz-Babilonia, C., Musharbash, F., & Jain, A. (2022). 100. Disparities in elective spine surgery for Medicaid beneficiaries: a systematic review. *The Spine Journal*, 22(9), S54-S55.
26. Groundland, J., Brown, J. M., Monument, M., Bernthal, N., Jones, K. B., & Randall, R. L. (2022). What Are the Long-term Surgical Outcomes of Compressive Endoprosthetic Osseointegration of the Femur with a Minimum 10-year Follow-up Period?. *Clinical Orthopaedics and Related Research®*, 480(3), 539-548.
27. Glicksman, C., McGuire, P., Kadin, M., Lawrence, M., Haws, M., Newby, J., ... & Wixtrom, R. (2022). Impact of capsulectomy type on post-explantation systemic symptom improvement: findings from the ASERF systemic symptoms in women-biospecimen analysis study: part 1. *Aesthetic Surgery Journal*, 42(7), 809-819.
28. Gramlich, Y., Hofmann, L., Kress, S., Ruckes, C., Kemmerer, M., Klug, A., ... & Kremer, M. (2022). Critically high metal ion levels found in metal-on-metal modular hinged knee arthroplasty: a comparison of two different systems. *The Bone & Joint Journal*, 104(3), 376-385.
29. Miséré, R. M., & van der Hulst, R. R. (2022). Self-reported health complaints in women undergoing explantation of breast implants. *Aesthetic Surgery Journal*, 42(2), 171-180.
30. Tan, E. S., Lee, J. Y., Boey, E., Soh, R., Sim, M. G., Yeo, W. T., ... & Kojodjojo, P. (2022). Use of extendable helix leads for conduction system pacing: Differences in lead handling and performance lead design impacts conduction system pacing. *Journal of Cardiovascular Electrophysiology*, 33(7), 1550-1557.
31. Patil, C., Agrawal, A., Abullais, S. S., Arora, S., Khateeb, S. U., & Fadul A. Elagib, M. (2022). Effectiveness of Different Chemotherapeutic Agents for Decontamination of Infected Dental Implant Surface: A Systematic Review. *Antibiotics*, 11(5), 593.

32. Zhu, W. Y., Guo, J., Yang, W. F., Tao, Z. Y., Lan, X., Wang, L., ... & Su, Y. X. (2022). Biodegradable magnesium implant enhances angiogenesis and alleviates medication-related osteonecrosis of the jaw in rats. *Journal of Orthopaedic Translation*, 33, 153-161.
- 33.